

**QK1 Rabbit mAb**  
**Catalog # AP75982****Specification****QK1 Rabbit mAb - Product Information**

Application	WB, IP
Primary Accession	<a href="#">Q96PU8</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	37671

**QK1 Rabbit mAb - Additional Information****Gene ID** 9444**Other Names**

QKI

**Dilution**

WB~~1/500-1/1000

IP~~1/20

**Format**

Liquid

**QK1 Rabbit mAb - Protein Information****Name** QKI {ECO:0000303|PubMed:16342280, ECO:0000312|HGNC:HGNC:21100}**Function**

RNA reader protein, which recognizes and binds specific RNAs, thereby regulating RNA metabolic processes, such as pre-mRNA splicing, circular RNA (circRNA) formation, mRNA export, mRNA stability and/or translation (PubMed:<a href="http://www.uniprot.org/citations/22398723" target="\_blank">22398723</a>, PubMed:<a href="http://www.uniprot.org/citations/23630077" target="\_blank">23630077</a>, PubMed:<a href="http://www.uniprot.org/citations/25768908" target="\_blank">25768908</a>, PubMed:<a href="http://www.uniprot.org/citations/27029405" target="\_blank">27029405</a>, PubMed:<a href="http://www.uniprot.org/citations/31331967" target="\_blank">31331967</a>, PubMed:<a href="http://www.uniprot.org/citations/37379838" target="\_blank">37379838</a>). Involved in various cellular processes, such as mRNA storage into stress granules, apoptosis, lipid deposition, interferon response, glial cell fate and development (PubMed:<a href="http://www.uniprot.org/citations/25768908" target="\_blank">25768908</a>, PubMed:<a href="http://www.uniprot.org/citations/31829086" target="\_blank">31829086</a>, PubMed:<a href="http://www.uniprot.org/citations/34428287" target="\_blank">34428287</a>, PubMed:<a href="http://www.uniprot.org/citations/37379838" target="\_blank">37379838</a>). Binds to the 5'-NACUAAAY-N(1,20)-UAAAY-3' RNA core sequence (PubMed:<a href="http://www.uniprot.org/citations/23630077" target="\_blank">23630077</a>). Acts as a mRNA modification reader that specifically recognizes and binds mRNA transcripts

modified by internal N(7)-methylguanine (m7G) (PubMed:<a href="http://www.uniprot.org/citations/37379838" target="\_blank">37379838</a>). Promotes the formation of circular RNAs (circRNAs) during the epithelial to mesenchymal transition and in cardiomyocytes: acts by binding to sites flanking circRNA-forming exons (PubMed:<a href="http://www.uniprot.org/citations/25768908" target="\_blank">25768908</a>). CircRNAs are produced by back-splicing circularization of pre-mRNAs (PubMed:<a href="http://www.uniprot.org/citations/25768908" target="\_blank">25768908</a>). Plays a central role in myelination via 3 distinct mechanisms (PubMed:<a href="http://www.uniprot.org/citations/16641098" target="\_blank">16641098</a>). First, acts by protecting and promoting stability of target mRNAs such as MBP, SIRT2 and CDKN1B, which promotes oligodendrocyte differentiation (By similarity). Second, participates in mRNA transport by regulating the nuclear export of MBP mRNA (By similarity). Finally, indirectly regulates mRNA splicing of MAG pre-mRNA during oligodendrocyte differentiation by acting as a negative regulator of MAG exon 12 alternative splicing: acts by binding to HNRNPA1 mRNA splicing factor, preventing its translation (By similarity). Involved in microglia differentiation and remyelination by regulating microexon alternative splicing of the Rho GTPase pathway (By similarity). Involved in macrophage differentiation: promotes monocyte differentiation by regulating pre-mRNA splicing in naive peripheral blood monocytes (PubMed:<a href="http://www.uniprot.org/citations/27029405" target="\_blank">27029405</a>). Acts as an important regulator of muscle development: required for the contractile function of cardiomyocytes by regulating alternative splicing of cardiomyocyte transcripts (By similarity). Acts as a negative regulator of thermogenesis by decreasing stability, nuclear export and translation of mRNAs encoding PPARGC1A and UCP1 (By similarity). Also required for visceral endoderm function and blood vessel development (By similarity). May also play a role in smooth muscle development (PubMed:<a href="http://www.uniprot.org/citations/31331967" target="\_blank">31331967</a>). In addition to its RNA-binding activity, also acts as a nuclear transcription coactivator for SREBF2/SREBP2 (By similarity).

#### **Cellular Location**

Nucleus. Cytoplasm [Isoform QKI6]: Cytoplasm, cytosol. Nucleus Note=Localizes predominantly in the cytoplasm and at lower levels in nucleus.

#### **Tissue Location**

Expressed in the frontal cortex of brain. Down-regulated in the brain of schizophrenic patients

#### **QK1 Rabbit mAb - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **QK1 Rabbit mAb - Images**



