

**Rab10 Rabbit mAb**  
**Catalog # AP75983****Specification**

---

**Rab10 Rabbit mAb - Product Information**

Application	WB, IP, ICC
Primary Accession	<a href="#">P61026</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	22541

**Rab10 Rabbit mAb - Additional Information****Gene ID** 10890**Other Names**  
RAB10**Dilution**  
WB~~1/500-1/1000  
IP~~N/A  
ICC~~N/A**Format**  
Liquid**Rab10 Rabbit mAb - Protein Information****Name** RAB10 ([HGNC:9759](#))**Function**

The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes (PubMed:<a href="http://www.uniprot.org/citations/21248164" target="\_blank">21248164</a>). Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different set of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion (PubMed:<a href="http://www.uniprot.org/citations/21248164" target="\_blank">21248164</a>). That Rab is mainly involved in the biosynthetic transport of proteins from the Golgi to the plasma membrane (PubMed:<a href="http://www.uniprot.org/citations/21248164" target="\_blank">21248164</a>). Regulates, for instance, SLC2A4/GLUT4 glucose transporter-enriched vesicles delivery to the plasma membrane (By similarity). In parallel, it regulates the transport of TLR4, a toll- like receptor to the plasma membrane and therefore may be important for innate immune response (By similarity). Also plays a specific role in asymmetric protein transport to the plasma membrane (PubMed:<a href="http://www.uniprot.org/citations/16641372" target="\_blank">16641372</a>). In neurons, it is involved in axonogenesis through regulation of vesicular membrane trafficking toward the axonal plasma membrane (By similarity). In epithelial cells, it regulates transport from the Golgi to

the basolateral membrane (PubMed:<a href="http://www.uniprot.org/citations/16641372" target="\_blank">16641372</a>). May play a role in the basolateral recycling pathway and in phagosome maturation (By similarity). May play a role in endoplasmic reticulum dynamics and morphology controlling tubulation along microtubules and tubules fusion (PubMed:<a href="http://www.uniprot.org/citations/23263280" target="\_blank">23263280</a>). Together with LRRK2, RAB8A, and RILPL1, it regulates ciliogenesis (PubMed:<a href="http://www.uniprot.org/citations/30398148" target="\_blank">30398148</a>). When phosphorylated by LRRK2 on Thr-73, binds RILPL1 and inhibits ciliogenesis (PubMed:<a href="http://www.uniprot.org/citations/30398148" target="\_blank">30398148</a>). Participates in the export of a subset of neosynthesized proteins through a Rab8- Rab10-Rab11-dependent endosomal export route (PubMed:<a href="http://www.uniprot.org/citations/32344433" target="\_blank">32344433</a>). Targeted to and stabilized on stressed lysosomes through LRRK2 phosphorylation where it promotes the extracellular release of lysosomal content through EHBP1 and EHNP1L1 effector proteins (PubMed:<a href="http://www.uniprot.org/citations/30209220" target="\_blank">30209220</a>).

### Cellular Location

Cytoplasmic vesicle membrane; Lipid-anchor; Cytoplasmic side. Golgi apparatus membrane. Golgi apparatus, trans-Golgi network membrane {ECO:0000250|UniProtKB:P24409}. Endosome membrane Recycling endosome membrane {ECO:0000250|UniProtKB:P24409}. Cytoplasmic vesicle, phagosome membrane {ECO:0000250|UniProtKB:P24409}. Cytoplasm, cytoskeleton, cilium basal body. Endoplasmic reticulum membrane. Cytoplasm, perinuclear region. Lysosome. Note=Associates with SLC2A4/GLUT4 storage vesicles (PubMed:22908308). Localizes to the base of the cilium when phosphorylated by LRRK2 on Thr-73 (PubMed:20576682, PubMed:30398148). Transiently associates with phagosomes (By similarity). Localizes to the endoplasmic reticulum at domains of new tubule growth (PubMed:23263280). Colocalizes with MICAL1, GRAF1/ARHGAP26 and GRAF2/ARHGAP10 on endosomal tubules (PubMed:32344433). Localizes to enlarged lysosomes through LRRK2 phosphorylation (PubMed:30209220). {ECO:0000250|UniProtKB:P24409, ECO:0000269|PubMed:20576682, ECO:0000269|PubMed:22908308, ECO:0000269|PubMed:23263280, ECO:0000269|PubMed:30209220, ECO:0000269|PubMed:30398148, ECO:0000269|PubMed:32344433}

### Tissue Location

Expressed in the hippocampus (PubMed:29562525). Expressed in neutrophils (at protein level) (PubMed:29127255) Expressed in the testis (at protein level) (PubMed:28067790)

## Rab10 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Rab10 Rabbit mAb - Images



