

**Rad51D Rabbit mAb**  
**Catalog # AP75994****Specification**

---

**Rad51D Rabbit mAb - Product Information**

Application	WB, IP
Primary Accession	<a href="#">O75771</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	35049

**Rad51D Rabbit mAb - Additional Information****Gene ID** 5892**Other Names**  
RAD51D**Dilution**  
WB~~1/500-1/1000  
IP~~N/A**Format**  
Liquid**Rad51D Rabbit mAb - Protein Information****Name** RAD51D**Synonyms** RAD51L3**Function**

Involved in the homologous recombination repair (HRR) pathway of double-stranded DNA breaks arising during DNA replication or induced by DNA-damaging agents. Bind to single-stranded DNA (ssDNA) and has DNA-dependent ATPase activity. Part of the RAD51 paralog protein complex BCDX2 which acts in the BRCA1-BRCA2-dependent HR pathway. Upon DNA damage, BCDX2 acts downstream of BRCA2 recruitment and upstream of RAD51 recruitment. BCDX2 binds predominantly to the intersection of the four duplex arms of the Holliday junction and to junction of replication forks. The BCDX2 complex was originally reported to bind single-stranded DNA, single-stranded gaps in duplex DNA and specifically to nicks in duplex DNA. Involved in telomere maintenance. The BCDX2 subcomplex XRCC2:RAD51D can stimulate Holliday junction resolution by BLM.

**Cellular Location**

Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Chromosome, telomere

**Tissue Location**

Expressed in colon, prostate, spleen, testis, ovary, thymus and small intestine. Weakly expressed in leukocytes

**Rad51D Rabbit mAb - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Rad51D Rabbit mAb - Images**