

**S100A4 Rabbit mAb**  
**Catalog # AP76046****Specification**

---

**S100A4 Rabbit mAb - Product Information**

Application	WB, IHC-P, IHC-F, IP, ICC
Primary Accession	<a href="#">P26447</a>
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	11729

**S100A4 Rabbit mAb - Additional Information****Gene ID** 6275**Other Names**  
S100A4**Dilution**  
WB~~1/500-1/1000  
IHC-P~~N/A  
IHC-F~~N/A  
IP~~N/A  
ICC~~N/A**Format**  
50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.**Storage**  
Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.**S100A4 Rabbit mAb - Protein Information****Name** S100A4**Synonyms** CAPL, MTS1**Function**  
Calcium-binding protein that plays a role in various cellular processes including motility, angiogenesis, cell differentiation, apoptosis, and autophagy (PubMed:<a href="http://www.uniprot.org/citations/16707441" target="\_blank">16707441</a>, PubMed:<a href="http://www.uniprot.org/citations/23752197" target="\_blank">23752197</a>, PubMed:<a href="http://www.uniprot.org/citations/30713770" target="\_blank">30713770</a>). Increases cell motility and invasiveness by interacting with non-muscle myosin heavy chain (NMMHC) IIA/MYH9 (PubMed:<a href="http://www.uniprot.org/citations/16707441" target="\_blank">16707441</a>). Mechanistically, promotes filament depolymerization and increases the amount of soluble myosin-IIA, resulting in the formation of stable protrusions

facilitating chemotaxis (By similarity). Also modulates the pro-apoptotic function of TP53 by binding to its C-terminal transactivation domain within the nucleus and reducing its protein levels (PubMed:<a href="http://www.uniprot.org/citations/23752197" target="\_blank">23752197</a>). Within the extracellular space, stimulates cytokine production including granulocyte colony-stimulating factor and CCL24 from T-lymphocytes (By similarity). In addition, stimulates T-lymphocyte chemotaxis by acting as a chemoattractant complex with PGLYRP1 that promotes lymphocyte migration via CCR5 and CXCR3 receptors (PubMed:<a href="http://www.uniprot.org/citations/26654597" target="\_blank">26654597</a>, PubMed:<a href="http://www.uniprot.org/citations/30713770" target="\_blank">30713770</a>).

#### Cellular Location

Secreted. Nucleus Cytoplasm {ECO:0000250|UniProtKB:P07091}

#### Tissue Location

Ubiquitously expressed.

### S100A4 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### S100A4 Rabbit mAb - Images



