

SESN2 Rabbit mAb
Catalog # AP76071**Specification****SESN2 Rabbit mAb - Product Information**

Application	WB, IP, ICC
Primary Accession	P58004
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	54494

SESN2 Rabbit mAb - Additional Information**Gene ID** 83667**Other Names**
SESN2**Dilution**
WB~~1/500-1/1000
IP~~1/20
ICC~~N/A**Format**
Liquid**SESN2 Rabbit mAb - Protein Information****Name** SESN2 ([HGNC:20746](#))**Function**

Functions as an intracellular leucine sensor that negatively regulates the mTORC1 signaling pathway through the GATOR complex (PubMed:18692468, PubMed:25263562, PubMed:25457612, PubMed:26449471, PubMed:26586190, PubMed:26612684, PubMed:31586034, PubMed:35114100, PubMed:35831510, PubMed:36528027). In absence of leucine, binds the GATOR subcomplex GATOR2 and prevents mTORC1 signaling (PubMed:18692468, PubMed:25263562, PubMed:25457612, PubMed:<a

[26449471](http://www.uniprot.org/citations/26449471), PubMed: [26586190](http://www.uniprot.org/citations/26586190), PubMed: [26612684](http://www.uniprot.org/citations/26612684), PubMed: [31586034](http://www.uniprot.org/citations/31586034), PubMed: [35114100](http://www.uniprot.org/citations/35114100), PubMed: [35831510](http://www.uniprot.org/citations/35831510), PubMed: [36528027](http://www.uniprot.org/citations/36528027)). Binding of leucine to SESN2 disrupts its interaction with GATOR2 thereby activating the TORC1 signaling pathway (PubMed: [26449471](http://www.uniprot.org/citations/26449471), PubMed: [26586190](http://www.uniprot.org/citations/26586190), PubMed: [35114100](http://www.uniprot.org/citations/35114100), PubMed: [35831510](http://www.uniprot.org/citations/35831510), PubMed: [36528027](http://www.uniprot.org/citations/36528027)). This stress-inducible metabolic regulator also plays a role in protection against oxidative and genotoxic stresses. May negatively regulate protein translation in response to endoplasmic reticulum stress, via mTORC1 (PubMed: [24947615](http://www.uniprot.org/citations/24947615)). May positively regulate the transcription by NFE2L2 of genes involved in the response to oxidative stress by facilitating the SQSTM1-mediated autophagic degradation of KEAP1 (PubMed: [23274085](http://www.uniprot.org/citations/23274085)). May also mediate TP53 inhibition of TORC1 signaling upon genotoxic stress (PubMed: [18692468](http://www.uniprot.org/citations/18692468)). Moreover, may prevent the accumulation of reactive oxygen species (ROS) through the alkylhydroperoxide reductase activity born by the N- terminal domain of the protein (PubMed: [26612684](http://www.uniprot.org/citations/26612684)). Was originally reported to contribute to oxidative stress resistance by reducing PRDX1 (PubMed: [15105503](http://www.uniprot.org/citations/15105503)). However, this could not be confirmed (PubMed: [19113821](http://www.uniprot.org/citations/19113821)).

Cellular Location

Cytoplasm.

Tissue Location

Widely expressed..

SESN2 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SESN2 Rabbit mAb - Images

