

SHP1 Rabbit mAb
Catalog # AP76080**Specification**

SHP1 Rabbit mAb - Product Information

Application	WB, IHC-P
Primary Accession	P29350
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	67561

SHP1 Rabbit mAb - Additional Information**Gene ID** 5777**Other Names**
PTPN6**Dilution**
WB~~1/500-1/1000
IHC-P~~N/A**Format**
50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.**Storage**
Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.**SHP1 Rabbit mAb - Protein Information****Name** PTPN6**Synonyms** HCP, PTP1C**Function**

Tyrosine phosphatase enzyme that plays important roles in controlling immune signaling pathways and fundamental physiological processes such as hematopoiesis (PubMed:14739280, PubMed:29925997). Dephosphorylates and negatively regulate several receptor tyrosine kinases (RTKs) such as EGFR, PDGFR and FGFR, thereby modulating their signaling activities (PubMed:21258366, PubMed:9733788). When recruited to immunoreceptor tyrosine-based inhibitory motif (ITIM)-containing receptors such as immunoglobulin-like transcript 2/LILRB1, programmed cell death protein 1/PDCD1, CD3D, CD22, CLEC12A and other receptors involved in immune regulation, initiates their dephosphorylation and subsequently inhibits downstream signaling events (PubMed:<a

[11907092](http://www.uniprot.org/citations/11907092), PubMed: [14739280](http://www.uniprot.org/citations/14739280), PubMed: [37932456](http://www.uniprot.org/citations/37932456), PubMed: [38166031](http://www.uniprot.org/citations/38166031)). Modulates the signaling of several cytokine receptors including IL-4 receptor (PubMed: [9065461](http://www.uniprot.org/citations/9065461)). Additionally, targets multiple cytoplasmic signaling molecules including STING1, LCK or STAT1 among others involved in diverse cellular processes including modulation of T-cell activation or cGAS-STING signaling (PubMed: [34811497](http://www.uniprot.org/citations/34811497), PubMed: [38532423](http://www.uniprot.org/citations/38532423)). Within the nucleus, negatively regulates the activity of some transcription factors such as NFAT5 via direct dephosphorylation. Also acts as a key transcriptional regulator of hepatic gluconeogenesis by controlling recruitment of RNA polymerase II to the PCK1 promoter together with STAT5A (PubMed: [37595871](http://www.uniprot.org/citations/37595871)).

Cellular Location

Cytoplasm. Nucleus Note=In neurons, translocates into the nucleus after treatment with angiotensin II (By similarity). Shuttles between the cytoplasm and nucleus via its association with PDPK1.

Tissue Location

Isoform 1 is expressed in hematopoietic cells. Isoform 2 is expressed in non-hematopoietic cells

SHP1 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SHP1 Rabbit mAb - Images



