

STK3 Rabbit mAb
Catalog # AP76127**Specification****STK3 Rabbit mAb - Product Information**

Application	WB, IHC-P, IP
Primary Accession	Q13188
Reactivity	Human, Mouse, Rat, Hamster
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	56301

STK3 Rabbit mAb - Additional Information**Gene ID** 6788**Other Names**
STK3**Dilution**
WB~~1/500-1/1000
IHC-P~~N/A
IP~~1/20**Format**
Liquid**STK3 Rabbit mAb - Protein Information****Name** STK3 ([HGNC:11406](#))**Function**

Stress-activated, pro-apoptotic kinase which, following caspase-cleavage, enters the nucleus and induces chromatin condensation followed by internucleosomal DNA fragmentation (PubMed:11278283, PubMed:8566796, PubMed:8816758). Key component of the Hippo signaling pathway which plays a pivotal role in organ size control and tumor suppression by restricting proliferation and promoting apoptosis. The core of this pathway is composed of a kinase cascade wherein STK3/MST2 and STK4/MST1, in complex with its regulatory protein SAV1, phosphorylates and activates LATS1/2 in complex with its regulatory protein MOB1, which in turn phosphorylates and inactivates YAP1 oncprotein and WWTR1/TAZ (PubMed:15688006, PubMed:16930133, PubMed:23972470, PubMed:28087714, PubMed:29063833, PubMed:30622739).

Phosphorylation of YAP1 by LATS2 inhibits its translocation into the nucleus to regulate cellular genes important for cell proliferation, cell death, and cell migration (PubMed:15688006, PubMed:16930133, PubMed:23972470, PubMed:28087714). STK3/MST2 and STK4/MST1 are required to repress proliferation of mature hepatocytes, to prevent activation of facultative adult liver stem cells (oval cells), and to inhibit tumor formation. Phosphorylates NKX2-1 (By similarity). Phosphorylates NEK2 and plays a role in centrosome disjunction by regulating the localization of NEK2 to centrosome, and its ability to phosphorylate CROCC and CEP250 (PubMed:21076410, PubMed:21723128). In conjunction with SAV1, activates the transcriptional activity of ESR1 through the modulation of its phosphorylation (PubMed:21104395). Positively regulates RAF1 activation via suppression of the inhibitory phosphorylation of RAF1 on 'Ser-259' (PubMed:20212043). Phosphorylates MOBKL1A and RASSF2 (PubMed:19525978). Phosphorylates MOBKL1B on 'Thr- 74'. Acts cooperatively with MOBKL1B to activate STK38 (PubMed:18328708, PubMed:18362890).

Cellular Location

Cytoplasm. Nucleus Cytoplasm, cytoskeleton, microtubule organizing center, centrosome.
Note=The caspase-cleaved form cycles between nucleus and cytoplasm (PubMed:11278283, PubMed:19525978) Phosphorylation at Thr-117 leads to inhibition of nuclear translocation (PubMed:19525978).

Tissue Location

Expressed at high levels in adult kidney, skeletal and placenta tissues and at very low levels in adult heart, lung and brain tissues.

STK3 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

STK3 Rabbit mAb - Images

