

**TRIB3 Rabbit mAb**  
**Catalog # AP76199****Specification**

---

**TRIB3 Rabbit mAb - Product Information**

Application	WB, IP
Primary Accession	<a href="#">Q96RU7</a>
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	39578

**TRIB3 Rabbit mAb - Additional Information****Gene ID** 57761**Other Names**  
TRIB3**Dilution**  
WB~~1/500-1/1000  
IP~~N/A**Format**  
50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.**Storage**  
Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.**TRIB3 Rabbit mAb - Protein Information****Name** TRIB3**Synonyms** C20orf97, NIPK, SKIP3, TRB3**Function**

Inactive protein kinase which acts as a regulator of the integrated stress response (ISR), a process for adaptation to various stress (PubMed:<a href="http://www.uniprot.org/citations/15775988" target="\_blank">15775988</a>, PubMed:<a href="http://www.uniprot.org/citations/15781252" target="\_blank">15781252</a>). Inhibits the transcriptional activity of DDIT3/CHOP and is involved in DDIT3/CHOP-dependent cell death during ER stress (PubMed:<a href="http://www.uniprot.org/citations/15775988" target="\_blank">15775988</a>, PubMed:<a href="http://www.uniprot.org/citations/15781252" target="\_blank">15781252</a>). May play a role in programmed neuronal cell death but does not appear to affect non-neuronal cells (PubMed:<a href="http://www.uniprot.org/citations/15775988" target="\_blank">15775988</a>, PubMed:<a href="http://www.uniprot.org/citations/15781252" target="\_blank">15781252</a>). Acts as a negative feedback regulator of the ATF4-dependent transcription during the ISR: while TRIB3 expression is promoted by ATF4, TRIB3 protein interacts with ATF4 and inhibits ATF4

transcription activity (By similarity). Disrupts insulin signaling by binding directly to Akt kinases and blocking their activation (By similarity). May bind directly to and mask the 'Thr-308' phosphorylation site in AKT1 (By similarity). Interacts with the NF-kappa-B transactivator p65 RELA and inhibits its phosphorylation and thus its transcriptional activation activity (PubMed:<a href="http://www.uniprot.org/citations/12736262" target="\_blank">12736262</a>). Interacts with MAPK kinases and regulates activation of MAP kinases (PubMed:<a href="http://www.uniprot.org/citations/15299019" target="\_blank">15299019</a>). Can inhibit APOBEC3A editing of nuclear DNA (PubMed:<a href="http://www.uniprot.org/citations/22977230" target="\_blank">22977230</a>).

### Cellular Location

Nucleus.

### Tissue Location

Highest expression in liver, pancreas, peripheral blood leukocytes and bone marrow. Also highly expressed in a number of primary lung, colon and breast tumors. Expressed in spleen, thymus, and prostate and is undetectable in other examined tissues, including testis, ovary, small intestine, colon, leukocyte, heart, brain, placenta, lung, skeletal muscle, and kidney

### TRIB3 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### TRIB3 Rabbit mAb - Images

