

TRIB3 Rabbit mAb
Catalog # AP76199**Specification**

TRIB3 Rabbit mAb - Product Information

Application	WB, IP
Primary Accession	Q96RU7
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	39578

TRIB3 Rabbit mAb - Additional Information**Gene ID** 57761**Other Names**
TRIB3**Dilution**
WB~~1/500-1/1000
IP~~N/A**Format**
Liquid**TRIB3 Rabbit mAb - Protein Information****Name** TRIB3**Synonyms** C20orf97, NIPK, SKIP3, TRB3**Function**

Inactive protein kinase which acts as a regulator of the integrated stress response (ISR), a process for adaptation to various stress (PubMed: [15775988](http://www.uniprot.org/citations/15775988), PubMed: [15781252](http://www.uniprot.org/citations/15781252)). Inhibits the transcriptional activity of DDIT3/CHOP and is involved in DDIT3/CHOP-dependent cell death during ER stress (PubMed: [15775988](http://www.uniprot.org/citations/15775988), PubMed: [15781252](http://www.uniprot.org/citations/15781252)). May play a role in programmed neuronal cell death but does not appear to affect non-neuronal cells (PubMed: [15775988](http://www.uniprot.org/citations/15775988), PubMed: [15781252](http://www.uniprot.org/citations/15781252)). Acts as a negative feedback regulator of the ATF4-dependent transcription during the ISR: while TRIB3 expression is promoted by ATF4, TRIB3 protein interacts with ATF4 and inhibits ATF4 transcription activity (By similarity). Disrupts insulin signaling by binding directly to Akt kinases and blocking their activation (By similarity). May bind directly to and mask the 'Thr-308' phosphorylation site in AKT1 (By similarity). Interacts with the NF-kappa-B transactivator p65 RELA

and inhibits its phosphorylation and thus its transcriptional activation activity (PubMed:12736262). Interacts with MAPK kinases and regulates activation of MAP kinases (PubMed:15299019). Can inhibit APOBEC3A editing of nuclear DNA (PubMed:22977230).

Cellular Location

Nucleus.

Tissue Location

Highest expression in liver, pancreas, peripheral blood leukocytes and bone marrow. Also highly expressed in a number of primary lung, colon and breast tumors. Expressed in spleen, thymus, and prostate and is undetectable in other examined tissues, including testis, ovary, small intestine, colon, leukocyte, heart, brain, placenta, lung, skeletal muscle, and kidney

TRIB3 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TRIB3 Rabbit mAb - Images

