

**TFG Rabbit mAb**  
**Catalog # AP76205****Specification**

---

**TFG Rabbit mAb - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB, IHC-P, IHC-F, ICC  |
| Primary Accession | <a href="#">Q92734</a> |
| Reactivity        | Human, Mouse, Rat      |
| Host              | Rabbit                 |
| Clonality         | Monoclonal Antibody    |
| Calculated MW     | 43448                  |

**TFG Rabbit mAb - Additional Information****Gene ID** 10342**Other Names**  
TFG**Dilution**  
WB~~1/500-1/1000  
IHC-P~~N/A  
IHC-F~~N/A  
ICC~~N/A**Format**  
50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.**Storage**  
Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.**TFG Rabbit mAb - Protein Information****Name** TFG**Function**  
Plays a role in the normal dynamic function of the endoplasmic reticulum (ER) and its associated microtubules (PubMed:<a href="http://www.uniprot.org/citations/23479643" target="\_blank">23479643</a>, PubMed:<a href="http://www.uniprot.org/citations/27813252" target="\_blank">27813252</a>). Required for secretory cargo traffic from the endoplasmic reticulum to the Golgi apparatus (PubMed:<a href="http://www.uniprot.org/citations/21478858" target="\_blank">21478858</a>).**Cellular Location**  
Endoplasmic reticulum. Note=Localizes to endoplasmic reticulum exit site (ERES), also known as transitional endoplasmic reticulum (tER) (PubMed:21478858, PubMed:27813252).**Tissue Location**

Ubiquitous.

### TFG Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### TFG Rabbit mAb - Images



