

**Transcription Termination Factor 2 Rabbit mAb**  
**Catalog # AP76214****Specification**

---

**Transcription Termination Factor 2 Rabbit mAb - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB, IHC-P, IHC-F, ICC  |
| Primary Accession | <a href="#">Q9UNY4</a> |
| Reactivity        | Human, Rat             |
| Host              | Rabbit                 |
| Clonality         | Monoclonal Antibody    |
| Calculated MW     | 129588                 |

**Transcription Termination Factor 2 Rabbit mAb - Additional Information****Gene ID** 8458**Other Names**  
TTF2**Dilution**  
WB~~1/500-1/1000  
IHC-P~~N/A  
IHC-F~~N/A  
ICC~~N/A**Format**  
Liquid**Transcription Termination Factor 2 Rabbit mAb - Protein Information****Name** TTF2**Function**  
DsDNA-dependent ATPase which acts as a transcription termination factor by coupling ATP hydrolysis with removal of RNA polymerase II from the DNA template. May contribute to mitotic transcription repression. May also be involved in pre-mRNA splicing.**Cellular Location**  
Cytoplasm. Nucleus. Note=Cytoplasmic during interphase Relocates to the nucleus as cells enter mitosis**Transcription Termination Factor 2 Rabbit mAb - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Transcription Termination Factor 2 Rabbit mAb - Images



