

UFD1 Rabbit mAb

Catalog # AP76230

Specification

UFD1 Rabbit mAb - Product Information

Application
Primary Accession
Reactivity
Host
Clonality

WB 092890 Human Rabbit

Monoclonal Antibody

34500

UFD1 Rabbit mAb - Additional Information

Gene ID 7353

Calculated MW

Other Names UFD1

Dilution

WB~~1/500-1/1000

Format

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.

Storage

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

UFD1 Rabbit mAb - Protein Information

Name UFD1 (HGNC:12520)

Synonyms UFD1L

Function

Essential component of the ubiquitin-dependent proteolytic pathway which degrades ubiquitin fusion proteins. The ternary complex containing UFD1, VCP and NPLOC4 binds ubiquitinated proteins and is necessary for the export of misfolded proteins from the ER to the cytoplasm, where they are degraded by the proteasome. The NPLOC4-UFD1- VCP complex regulates spindle disassembly at the end of mitosis and is necessary for the formation of a closed nuclear envelope. It may be involved in the development of some ectoderm-derived structures (By similarity). Acts as a negative regulator of type I interferon production via the complex formed with VCP and NPLOC4, which binds to RIGI and recruits RNF125 to promote ubiquitination and degradation of RIGI (PubMed:26471729).

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q9ES53}. Cytoplasm, cytosol





Tel: 858.875.1900 Fax: 858.875.1999

{ECO:0000250|UniProtKB:Q9ES53}

Tissue Location

Found in adult heart, skeletal muscle and pancreas, and in fetal liver and kidney

UFD1 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

UFD1 Rabbit mAb - Images

