

Wnt5a Rabbit mAb

Catalog # AP76261

Specification

Wnt5a Rabbit mAb - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW

WB, ICC
P41221
Human
Rabbit
Monoclonal Antibody

42339

Wnt5a Rabbit mAb - Additional Information

Gene ID 7474

Other Names WNT5A

DilutionWB~~1/500-1/1000
ICC~~N/A

Format Liquid

Wnt5a Rabbit mAb - Protein Information

Name WNT5A

Function

Ligand for members of the frizzled family of seven transmembrane receptors. Can activate or inhibit canonical Wnt signaling, depending on receptor context. In the presence of FZD4, activates beta-catenin signaling. In the presence of ROR2, inhibits the canonical Wnt pathway by promoting beta-catenin degradation through a GSK3-independent pathway which involves down-regulation of beta- catenin-induced reporter gene expression (By similarity). Suppression of the canonical pathway allows chondrogenesis to occur and inhibits tumor formation. Stimulates cell migration. Decreases proliferation, migration, invasiveness and clonogenicity of carcinoma cells and may act as a tumor suppressor (PubMed:15735754 (PubMed:17426020 (PubMed:<a href="http://www.uniprot.org/citations/17426020" target="_blank") (PubMed:<a href="http://www.unip

Cellular Location

Secreted, extracellular space, extracellular matrix. Secreted

Tissue Location





Tel: 858.875.1900 Fax: 858.875.1999

Expression is increased in differentiated thyroid carcinomas compared to normal thyroid tissue and anaplastic thyroid tumors where expression is low or undetectable. Expression is found in thyrocytes but not in stromal cells (at protein level) (PubMed:15735754). Detected in neonate heart and lung (PubMed:8288227)

Wnt5a Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Wnt5a Rabbit mAb - Images







