

BAP1 Rabbit mAb
Catalog # AP76400**Specification****BAP1 Rabbit mAb - Product Information**

Application	WB
Primary Accession	Q92560
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	80362

BAP1 Rabbit mAb - Additional Information

Gene ID 8314

Other Names

BAP1

Dilution

WB~~1/500-1/1000

Format

Liquid

BAP1 Rabbit mAb - Protein Information**Name** BAP1 {ECO:0000303|PubMed:9528852, ECO:0000312|HGNC:HGNC:950}**Function**

Deubiquitinating enzyme that plays a key role in chromatin by mediating deubiquitination of histone H2A and HCFC1 (PubMed: [12485996](http://www.uniprot.org/citations/12485996), PubMed: [18757409](http://www.uniprot.org/citations/18757409), PubMed: [20436459](http://www.uniprot.org/citations/20436459), PubMed: [25451922](http://www.uniprot.org/citations/25451922), PubMed: [35051358](http://www.uniprot.org/citations/35051358)). Catalytic component of the polycomb repressive deubiquitinase (PR-DUB) complex, a complex that specifically mediates deubiquitination of histone H2A monoubiquitinated at 'Lys-120' (H2AK119ub1) (PubMed: [20436459](http://www.uniprot.org/citations/20436459), PubMed: [25451922](http://www.uniprot.org/citations/25451922), PubMed: [30664650](http://www.uniprot.org/citations/30664650), PubMed: [35051358](http://www.uniprot.org/citations/35051358)). Does not deubiquitinate monoubiquitinated histone H2B (PubMed: [20436459](http://www.uniprot.org/citations/20436459), PubMed: [30664650](http://www.uniprot.org/citations/30664650)). The PR-DUB complex is an epigenetic regulator of gene expression and acts as a transcriptional coactivator, affecting genes involved in development, cell communication, signaling, cell proliferation and cell

viability (PubMed:20805357, PubMed:30664650, PubMed:36180891). Antagonizes PRC1 mediated H2AK119ub1 monoubiquitination (PubMed:30664650). As part of the PR-DUB complex, associates with chromatin enriched in histone marks H3K4me1, H3K4me3, and H3K27Ac, but not in H3K27me3 (PubMed:36180891). Recruited to specific gene-regulatory regions by YY1 (PubMed:20805357). Acts as a regulator of cell growth by mediating deubiquitination of HCFC1 N- terminal and C-terminal chains, with some specificity toward 'Lys-48'- linked polyubiquitin chains compared to 'Lys-63'-linked polyubiquitin chains (PubMed:19188440, PubMed:19815555). Deubiquitination of HCFC1 does not lead to increase stability of HCFC1 (PubMed:19188440, PubMed:19815555). Interferes with the BRCA1 and BARD1 heterodimer activity by inhibiting their ability to mediate ubiquitination and autoubiquitination (PubMed:19117993). It however does not mediate deubiquitination of BRCA1 and BARD1 (PubMed:19117993). Able to mediate autodeubiquitination via intramolecular interactions to counteract monoubiquitination at the nuclear localization signal (NLS), thereby protecting it from cytoplasmic sequestration (PubMed:24703950). Negatively regulates epithelial-mesenchymal transition (EMT) of trophoblast stem cells during placental development by regulating genes involved in epithelial cell integrity, cell adhesion and cytoskeletal organization (PubMed:34170818).

Cellular Location

Cytoplasm. Nucleus. Chromosome. Note=Mainly nuclear (PubMed:24703950, PubMed:30664650). Binds to chromatin (PubMed:30664650). Localizes to the cytoplasm when monoubiquitinated by the E2/E3 hybrid ubiquitin- protein ligase UBE2O (PubMed:24703950). Recruitment to chromatin is dependent on ASXL1/2/3 and recruitment to specific genes on FOXK1/2 (By similarity). Nuclear localization is redundantly mediated by the importin and transportin systems; TNPO1/transportin-1 is the major mediator of nuclear localization (PubMed:35446349)
{ECO:0000250|UniProtKB:Q99PU7, ECO:0000269|PubMed:24703950, ECO:0000269|PubMed:30664650, ECO:0000269|PubMed:35446349}

Tissue Location

Highly expressed in testis, placenta and ovary (PubMed:9528852). Expressed in breast (PubMed:9528852). levels in the placenta increase over the course of pregnancy (PubMed:34170818)

BAP1 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

BAP1 Rabbit mAb - Images

