

Cullin 4A Rabbit mAb
Catalog # AP76456**Specification****Cullin 4A Rabbit mAb - Product Information**

Application	WB, IHC-P, IP
Primary Accession	Q13619
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	87680

Cullin 4A Rabbit mAb - Additional Information**Gene ID** 8451**Other Names**
CUL4A**Dilution**WB~~1/500-1/1000
IHC-P~~N/A
IP~~N/A**Format**
Liquid**Cullin 4A Rabbit mAb - Protein Information****Name** CUL4A {ECO:0000303|PubMed:9721878, ECO:0000312|HGNC:HGNC:2554}**Function**

Core component of multiple cullin-RING-based E3 ubiquitin-protein ligase complexes which mediate the ubiquitination of target proteins (PubMed:14578910, PubMed:14739464, PubMed:15448697, PubMed:15548678, PubMed:15811626, PubMed:16678110, PubMed:17041588, PubMed:24209620, PubMed:30166453, PubMed:33854232, PubMed:33854239). As a scaffold protein may contribute to catalysis through positioning of the substrate and the ubiquitin-conjugating enzyme (PubMed:14578910, PubMed:<a href="http://www.uniprot.org/citations/14739464"

target="_blank">>14739464, PubMed:>15448697, PubMed:>15548678, PubMed:>15811626, PubMed:>16678110, PubMed:>17041588, PubMed:>24209620). The E3 ubiquitin-protein ligase activity of the complex is dependent on the neddylation of the cullin subunit and is inhibited by the association of the deneddylated cullin subunit with TIP120A/CAND1 (PubMed:>14578910, PubMed:>14739464, PubMed:>15448697, PubMed:>15548678, PubMed:>15811626, PubMed:>16678110, PubMed:>17041588, PubMed:>24209620). The functional specificity of the E3 ubiquitin-protein ligase complex depends on the variable substrate recognition component (PubMed:>14578910, PubMed:>14739464, PubMed:>15448697, PubMed:>15548678, PubMed:>15811626, PubMed:>16678110, PubMed:>17041588, PubMed:>24209620). DCX(DET1-COP1) directs ubiquitination of JUN (PubMed:>14739464). DCX(DDB2) directs ubiquitination of XPC (PubMed:>15811626). DCX(DDB2) ubiquitinates histones H3-H4 and is required for efficient histone deposition during replication-coupled (H3.1) and replication-independent (H3.3) nucleosome assembly, probably by facilitating the transfer of H3 from ASF1A/ASF1B to other chaperones involved in histone deposition (PubMed:>16678110, PubMed:>17041588, PubMed:>24209620). DCX(DTL) plays a role in PCNA-dependent polyubiquitination of CDT1 and MDM2-dependent ubiquitination of p53/TP53 in response to radiation-induced DNA damage and during DNA replication (PubMed:>14578910, PubMed:>15448697, PubMed:>15548678). DCX(DTL) directs autoubiquitination of DTL (PubMed:>23478445). In association with DDB1 and SKP2 probably is involved in ubiquitination of CDKN1B/p27kip (PubMed:>16537899). Is involved in ubiquitination of HOXA9 (PubMed:>14609952). The DDB1-CUL4A-DTL E3 ligase complex regulates the circadian clock function by mediating the ubiquitination and degradation of CRY1 (PubMed:>26431207). The DCX(ERCC8) complex (also named CSA complex) plays a role in transcription-coupled repair (TCR) (PubMed:>12732143, PubMed:>32355176, PubMed:>38316879). A number of DCX complexes (containing either TRPC4AP or DCAF12 as substrate-recognition component) are part of the DesCEND (destruction via C-end degrons) pathway, which recognizes a C-degron located at the extreme C terminus of target proteins, leading to their ubiquitination and degradation (PubMed:>29779948).

target="_blank">>29779948). The DCX(AMBRA1) complex is a master regulator of the transition from G1 to S cell phase by mediating ubiquitination of phosphorylated cyclin-D (CCND1, CCND2 and CCND3) (PubMed:33854232, PubMed:33854239). The DCX(AMBRA1) complex also acts as a regulator of Cul5-RING (CRL5) E3 ubiquitin-protein ligase complexes by mediating ubiquitination and degradation of Elongin-C (ELOC) component of CRL5 complexes (PubMed:30166453). With CUL4B, contributes to ribosome biogenesis (PubMed:26711351).

Cullin 4A Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Cullin 4A Rabbit mAb - Images



