

Dnmt3a Rabbit mAb

Catalog # AP76472

Specification

Dnmt3a Rabbit mAb - Product Information

Application WB
Primary Accession Q9Y6K1
Reactivity Human
Host Rabbit
Clonality Monoclonal Antibody

Calculated MW 101858

Dnmt3a Rabbit mAb - Additional Information

Gene ID 1788

Other Names DNMT3A

Dilution WB~~1/500-1/1000

Format Liquid

Dnmt3a Rabbit mAb - Protein Information

Name DNMT3A

Function

Required for genome-wide de novo methylation and is essential for the establishment of DNA methylation patterns during development (PubMed:<a

 $href="http://www.uniprot.org/citations/12138111" target="_blank">12138111, PubMed:16357870, PubMed:30478443). DNA methylation is coordinated with methylation of histones (PubMed:<a$

 $href="http://www.uniprot.org/citations/12138111" target="_blank">12138111, PubMed:16357870, PubMed:30478443), It modifies DNA in a non-processive manner and also methylates non-CpG sites (PubMed:<a$

href="http://www.uniprot.org/citations/12138111" target="_blank">12138111, PubMed:16357870, PubMed:30478443). May preferentially methylate DNA linker between 2 nucleosomal cores and is inhibited by histone H1 (By similarity). Plays a role in paternal and maternal imprinting (By similarity). Required for methylation of most imprinted loci in germ cells (By similarity). Acts as a transcriptional corepressor for ZBTB18 (By similarity). Recruited to trimethylated 'Lys-36' of histone H3 (H3K36me3) sites (By similarity). Can actively repress transcription through the recruitment of





HDAC activity (By similarity). Also has weak auto-methylation activity on Cys-710 in absence of DNA (By similarity).

Cellular Location

Nucleus. Chromosome Cytoplasm. Note=Accumulates in the major satellite repeats at pericentric heterochromatin {ECO:0000250|UniProtKB:088508}

Tissue Location

Highly expressed in fetal tissues, skeletal muscle, heart, peripheral blood mononuclear cells, kidney, and at lower levels in placenta, brain, liver, colon, spleen, small intestine and lung

Dnmt3a Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Dnmt3a Rabbit mAb - Images

