

**Sestrin-1 Antibody (N-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP7650a****Specification**

---

**Sestrin-1 Antibody (N-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q9Y6P5</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	14-43

**Sestrin-1 Antibody (N-term) - Additional Information****Gene ID** 27244**Other Names**

Sestrin-1, p53-regulated protein PA26, SESN1, PA26, SEST1

**Target/Specificity**

This Sestrin-1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 14-43 amino acids from the N-terminal region of human Sestrin-1.

**Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Sestrin-1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**Sestrin-1 Antibody (N-term) - Protein Information****Name** SESN1 ([HGNC:21595](#))

**Function** Functions as an intracellular leucine sensor that negatively regulates the TORC1 signaling pathway through the GATOR complex. In absence of leucine, binds the GATOR subcomplex GATOR2 and prevents TORC1 signaling. Binding of leucine to SESN2 disrupts its

interaction with GATOR2 thereby activating the TORC1 signaling pathway (PubMed:[25263562](#), PubMed:[26449471](#)). This stress-inducible metabolic regulator may also play a role in protection against oxidative and genotoxic stresses (By similarity). May positively regulate the transcription by NFE2L2 of genes involved in the response to oxidative stress by facilitating the SQSTM1-mediated autophagic degradation of KEAP1 (PubMed:[23274085](#)). Moreover, may prevent the accumulation of reactive oxygen species (ROS) through the alkylhydroperoxide reductase activity born by the N-terminal domain of the protein (By similarity). Was originally reported to contribute to oxidative stress resistance by reducing PRDX1 (PubMed:[15105503](#)). However, this could not be confirmed (By similarity).

**Cellular Location**

Nucleus. Cytoplasm


**Tissue Location**

Widely expressed..

**Sestrin-1 Antibody (N-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Sestrin-1 Antibody (N-term) - Images**

Western blot analysis of Sestrin-1 Antibody (N-term) (Cat. #AP7650a) in mouse heart tissue lysates (35ug/lane). Sestrin-1 (arrow) was detected using the purified Pab.

**Sestrin-1 Antibody (N-term) - Background**

Sestrin-1 is involved in the reduction of peroxiredoxins. This protein may also be regulator of cellular growth.

**Sestrin-1 Antibody (N-term) - References**

Budanov,A.V., Science 304 (5670), 596-600 (2004) Peeters,H., Hum. Genet. 112 (5-6), 573-580 (2003) Velasco-Miguel,S., Oncogene 18 (1), 127-137 (1999)

**Sestrin-1 Antibody (N-term) - Citations**

- [GCN2 sustains mTORC1 suppression upon amino acid deprivation by inducing Sestrin2.](#)