

Ku70 Rabbit mAb
Catalog # AP76564**Specification**

Ku70 Rabbit mAb - Product Information

| | |
|-------------------|------------------------|
| Application | WB, IHC-P, IHC-F, ICC |
| Primary Accession | P12956 |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | Monoclonal Antibody |
| Calculated MW | 69843 |

Ku70 Rabbit mAb - Additional Information**Gene ID** 2547**Other Names**
XRCC6**Dilution**

WB~~1/500-1/1000
IHC-P~~N/A
IHC-F~~N/A
ICC~~N/A

Format

Liquid

Ku70 Rabbit mAb - Protein Information**Name** XRCC6**Synonyms** G22P1**Function**

Single-stranded DNA-dependent ATP-dependent helicase that plays a key role in DNA non-homologous end joining (NHEJ) by recruiting DNA-PK to DNA (PubMed:11493912, PubMed:12145306, PubMed:20493174, PubMed:2466842, PubMed:7957065, PubMed:8621488, PubMed:9742108). Required for double-strand break repair and V(D)J recombination (PubMed:11493912, PubMed:12145306, PubMed:20493174, PubMed:>2466842, PubMed:>7957065, PubMed:>8621488, PubMed:>9742108). Also has a role in chromosome translocation (PubMed:>11493912, PubMed:>12145306, PubMed:>20493174, PubMed:>2466842, PubMed:>7957065, PubMed:>8621488, PubMed:>9742108). Has a role in chromosome translocation (PubMed:>11493912, PubMed:>12145306, PubMed:>20493174, PubMed:>2466842, PubMed:>7957065, PubMed:>8621488, PubMed:>9742108). The DNA helicase II complex binds preferentially to fork-like ends of double-stranded DNA in a cell cycle-dependent manner (PubMed:>11493912, PubMed:>12145306, PubMed:>20493174, PubMed:>2466842, PubMed:>7957065, PubMed:>8621488, PubMed:>9742108). It works in the 3'-5' direction (PubMed:>11493912, PubMed:>12145306, PubMed:>20493174, PubMed:>2466842, PubMed:>7957065, PubMed:>8621488, PubMed:>9742108). During NHEJ, the XRCC5-XRRC6 dimer performs the recognition step: it recognizes and binds to the broken ends of the DNA and protects them from further resection (PubMed:>11493912, PubMed:>12145306, PubMed:>20493174, PubMed:>2466842, PubMed:>7957065, PubMed:>8621488, PubMed:>9742108). Binding to DNA may be mediated by XRCC6 (PubMed:>11493912, PubMed:>12145306, PubMed:>20493174, PubMed:>2466842, PubMed:>7957065, PubMed:>8621488, PubMed:>9742108). The XRCC5-XRRC6 dimer acts as a regulatory subunit of the DNA-dependent protein kinase complex DNA-PK by increasing the affinity of the catalytic subunit PRKDC to DNA by 100-fold (PubMed:>11493912, PubMed:>12145306, PubMed:>20493174, PubMed:>2466842).

target="_blank">>2466842, PubMed:>7957065, PubMed:>8621488, PubMed:>9742108). The XRCC5-XRRC6 dimer is probably involved in stabilizing broken DNA ends and bringing them together (PubMed:>11493912, PubMed:>12145306, PubMed:>20493174, PubMed:>2466842, PubMed:>7957065, PubMed:>8621488, PubMed:>9742108). The assembly of the DNA-PK complex to DNA ends is required for the NHEJ ligation step (PubMed:>11493912, PubMed:>12145306, PubMed:>20493174, PubMed:>2466842, PubMed:>7957065, PubMed:>8621488, PubMed:>9742108). Probably also acts as a 5'-deoxyribose-5-phosphate lyase (5'-dRP lyase), by catalyzing the beta-elimination of the 5' deoxyribose-5-phosphate at an abasic site near double-strand breaks (PubMed:>20383123). 5'-dRP lyase activity allows to 'clean' the termini of abasic sites, a class of nucleotide damage commonly associated with strand breaks, before such broken ends can be joined (PubMed:>20383123). The XRCC5-XRRC6 dimer together with APEX1 acts as a negative regulator of transcription (PubMed:>8621488). In association with NAA15, the XRCC5-XRRC6 dimer binds to the osteocalcin promoter and activates osteocalcin expression (PubMed:>12145306). Plays a role in the regulation of DNA virus-mediated innate immune response by assembling into the HDP-RNP complex, a complex that serves as a platform for IRF3 phosphorylation and subsequent innate immune response activation through the cGAS-STING pathway (PubMed:>28712728). Negatively regulates apoptosis by interacting with BAX and sequestering it from the mitochondria (PubMed:>15023334). Might have deubiquitination activity, acting on BAX (PubMed:>18362350).

Cellular Location

Nucleus. Chromosome. Cytoplasm. Note=When trimethylated, localizes in the cytoplasm.

Ku70 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Ku70 Rabbit mAb - Images