

**NEK6 Rabbit mAb**  
Catalog # AP76614**Specification**

---

**NEK6 Rabbit mAb - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">O9HC98</a>
Reactivity	<b>Human, Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Monoclonal Antibody</b>
Calculated MW	<b>35714</b>

**NEK6 Rabbit mAb - Additional Information****Gene ID** 10783**Other Names**  
NEK6**Dilution**  
WB~~1/500-1/1000**Format**  
50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.**Storage**  
Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.**NEK6 Rabbit mAb - Protein Information****Name** NEK6 ([HGNC:7749](#))**Function**

Protein kinase which plays an important role in mitotic cell cycle progression (PubMed:[11516946](http://www.uniprot.org/citations/11516946)), PubMed:[14563848](http://www.uniprot.org/citations/14563848)). Required for chromosome segregation at metaphase-anaphase transition, robust mitotic spindle formation and cytokinesis (PubMed:[19414596](http://www.uniprot.org/citations/19414596)). Phosphorylates ATF4, CIRS1, PTN, RAD26L, RBBP6, RPS7, RPS6KB1, TRIP4, STAT3 and histones H1 and H3 (PubMed:[12054534](http://www.uniprot.org/citations/12054534)), PubMed:[20873783](http://www.uniprot.org/citations/20873783)). Phosphorylates KIF11 to promote mitotic spindle formation (PubMed:[19001501](http://www.uniprot.org/citations/19001501)). Involved in G2/M phase cell cycle arrest induced by DNA damage (PubMed:[18728393](http://www.uniprot.org/citations/18728393)). Inhibition of activity results in apoptosis. May contribute to tumorigenesis by suppressing p53/TP53-induced cancer cell senescence (PubMed:[21099361](http://www.uniprot.org/citations/21099361))

target="\_blank">21099361</a>). Phosphorylates EML4 at 'Ser-144', promoting its dissociation from microtubules during mitosis which is required for efficient chromosome congression (PubMed:<a href="http://www.uniprot.org/citations/31409757" target="\_blank">31409757</a>).

#### Cellular Location

Cytoplasm. Nucleus. Nucleus speckle. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle pole. Note=Colocalizes with APBB1 at the nuclear speckles. Colocalizes with PIN1 in the nucleus. Colocalizes with ATF4, CIRSR, ARHGAP33, ANKRA2, CDC42, NEK9, RAD26L, RBBP6, RPS7, TRIP4, RELB and PHF1 in the centrosome. Localizes to spindle microtubules in metaphase and anaphase and to the midbody during cytokinesis

#### Tissue Location

Ubiquitous, with highest expression in heart and skeletal muscle.

### NEK6 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### NEK6 Rabbit mAb - Images

