

SBDS Rabbit mAb
Catalog # AP76704**Specification**

SBDS Rabbit mAb - Product Information

Application	WB, IP, ICC
Primary Accession	Q9Y3A5
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	28764

SBDS Rabbit mAb - Additional Information**Gene ID** 51119**Other Names**
SBDS**Dilution**
WB~~1/500-1/1000
IP~~N/A
ICC~~N/A**Format**
Liquid**SBDS Rabbit mAb - Protein Information****Name** SBDS**Function**

Required for the assembly of mature ribosomes and ribosome biogenesis. Together with EFL1, triggers the GTP-dependent release of EIF6 from 60S pre-ribosomes in the cytoplasm, thereby activating ribosomes for translation competence by allowing 80S ribosome assembly and facilitating EIF6 recycling to the nucleus, where it is required for 60S rRNA processing and nuclear export. Required for normal levels of protein synthesis. May play a role in cellular stress resistance. May play a role in cellular response to DNA damage. May play a role in cell proliferation.

Cellular Location

Cytoplasm. Nucleus, nucleolus. Nucleus, nucleoplasm. Cytoplasm, cytoskeleton, spindle. Note=Primarily detected in the cytoplasm, and at low levels in nucleus and nucleolus (PubMed:17475909, PubMed:19602484). Detected in the nucleolus during G1 and G2 phase of the cell cycle, and diffusely distributed in the nucleus during S phase. Detected at the mitotic spindle. Colocalizes with the microtubule organizing center during interphase (PubMed:19759903).

Tissue Location

Widely expressed.

SBDS Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SBDS Rabbit mAb - Images



