

PRKCDBP Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7787c

Specification

PRKCDBP Antibody (Center) - Product Information

Application	WB,E
Primary Accession	O969G5
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	27701
Antigen Region	113-144

PRKCDBP Antibody (Center) - Additional Information

Gene ID 112464

Other Names

Protein kinase C delta-binding protein, Cavin-3, Serum deprivation response factor-related gene product that binds to C-kinase, hSRBC, PRKCDBP, SRBC

Target/Specificity

This PRKCDBP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 113-144 amino acids from the Central region of human PRKCDBP.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PRKCDBP Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

PRKCDBP Antibody (Center) - Protein Information

Name CAVIN3 ([HGNC:9400](#))

Synonyms PRKCDBP, SRBC

Function Regulates the traffic and/or budding of caveolae (PubMed:[19262564](#)). Plays a role in caveola formation in a tissue- specific manner. Required for the formation of caveolae in smooth muscle but not in the lung and heart endothelial cells. Regulates the equilibrium between cell surface-associated and cell surface- dissociated caveolae by promoting the rapid release of caveolae from the cell surface. Plays a role in the regulation of the circadian clock. Modulates the period length and phase of circadian gene expression and also regulates expression and interaction of the core clock components PER1/2 and CRY1/2 (By similarity).

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q91VJ2}. Membrane, caveola. Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q91VJ2}. Note=Localizes in the caveolae in a caveolin-dependent manner.

Tissue Location


Skeletal muscle, liver, stomach, lung, kidney and heart (at protein level). Strongly expressed in mammary and epithelial cells.

PRKCDBP Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PRKCDBP Antibody (Center) - Images



Western blot analysis of anti-PRKCDBP Antibody (Center) (Cat.#AP7787c) in A2058 cell line lysates (35ug/lane). PRKCDBP (arrow) was detected using the purified Pab.



Western blot analysis of PRKCDBP (arrow) using rabbit polyclonal PRKCDBP Antibody (Center) (Cat. #AP7787c). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the PRKCDBP gene (Lane 2) (Origene Technologies).

PRKCDBP Antibody (Center) - Background

PRKCDBP was identified as a binding protein of the protein kinase C, delta (PRKCD). The expression of this protein in cultured cell lines is strongly induced by serum starvation, and was found to be down-regulated in various cancer cell lines, suggesting the possible tumor suppressor function of this protein.

PRKCDBP Antibody (Center) - References

- Lee,J.H., Int. J. Cancer 122 (7), 1573-1584 (2008)
- Zochbauer-Muller,S., Oncogene 24 (41), 6249-6255 (2005)
- Xu,X.L., Cancer Res. 61 (21), 7943-7949 (2001)