

**EIF4A3 Rabbit mAb**  
**Catalog # AP77978****Specification****EIF4A3 Rabbit mAb - Product Information**

Application	WB
Primary Accession	<a href="#">P38919</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	46871

**EIF4A3 Rabbit mAb - Additional Information****Gene ID** 9775**Other Names**  
EIF4A3**Dilution**  
WB~~1/500-1/1000**Format**  
Liquid**EIF4A3 Rabbit mAb - Protein Information****Name** EIF4A3**Synonyms** DDX48, KIAA0111**Function**

ATP-dependent RNA helicase (PubMed:<a href="http://www.uniprot.org/citations/16170325" target="\_blank">16170325</a>). Involved in pre-mRNA splicing as component of the spliceosome (PubMed:<a href="http://www.uniprot.org/citations/11991638" target="\_blank">11991638</a>, PubMed:<a href="http://www.uniprot.org/citations/22961380" target="\_blank">22961380</a>, PubMed:<a href="http://www.uniprot.org/citations/28076346" target="\_blank">28076346</a>, PubMed:<a href="http://www.uniprot.org/citations/28502770" target="\_blank">28502770</a>, PubMed:<a href="http://www.uniprot.org/citations/29301961" target="\_blank">29301961</a>). Core component of the splicing-dependent multiprotein exon junction complex (EJC) deposited at splice junctions on mRNAs (PubMed:<a href="http://www.uniprot.org/citations/16170325" target="\_blank">16170325</a>, PubMed:<a href="http://www.uniprot.org/citations/16209946" target="\_blank">16209946</a>, PubMed:<a href="http://www.uniprot.org/citations/16314458" target="\_blank">16314458</a>, PubMed:<a href="http://www.uniprot.org/citations/16923391" target="\_blank">16923391</a>, PubMed:<a href="http://www.uniprot.org/citations/16931718" target="\_blank">16931718</a>, PubMed:<a href="http://www.uniprot.org/citations/19033377" target="\_blank">19033377</a>, PubMed:<a href="http://www.uniprot.org/citations/20479275" target="\_blank">20479275</a>). The EJC is a dynamic structure consisting of core proteins and

several peripheral nuclear and cytoplasmic associated factors that join the complex only transiently either during EJC assembly or during subsequent mRNA metabolism. The EJC marks the position of the exon-exon junction in the mature mRNA for the gene expression machinery and the core components remain bound to spliced mRNAs throughout all stages of mRNA metabolism thereby influencing downstream processes including nuclear mRNA export, subcellular mRNA localization, translation efficiency and nonsense-mediated mRNA decay (NMD). Its RNA-dependent ATPase and RNA-helicase activities are induced by CASC3, but abolished in presence of the MAGOH-RBM8A heterodimer, thereby trapping the ATP-bound EJC core onto spliced mRNA in a stable conformation. The inhibition of ATPase activity by the MAGOH-RBM8A heterodimer increases the RNA-binding affinity of the EJC. Involved in translational enhancement of spliced mRNAs after formation of the 80S ribosome complex. Binds spliced mRNA in sequence-independent manner, 20-24 nucleotides upstream of mRNA exon-exon junctions. Shows higher affinity for single-stranded RNA in an ATP-bound core EJC complex than after the ATP is hydrolyzed. Involved in the splicing modulation of BCL2L1/Bcl-X (and probably other apoptotic genes); specifically inhibits formation of proapoptotic isoforms such as Bcl-X(S); the function is different from the established EJC assembly (PubMed:<a href="http://www.uniprot.org/citations/22203037" target="\_blank">22203037</a>). Involved in craniofacial development (PubMed:<a href="http://www.uniprot.org/citations/24360810" target="\_blank">24360810</a>).

#### Cellular Location

Nucleus. Nucleus speckle. Cytoplasm {ECO:0000250|UniProtKB:Q3B8Q2}.

Note=Nucleocytoplasmic shuttling protein. Travels to the cytoplasm as part of the exon junction complex (EJC) bound to mRNA. Detected in dendritic layer as well as the nuclear and cytoplasmic (somatic) compartments of neurons. Colocalizes with STAU1 and FMR1 in dendrites (By similarity) {ECO:0000250|UniProtKB:Q3B8Q2}

#### Tissue Location

Ubiquitously expressed.

### eIF4A3 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### eIF4A3 Rabbit mAb - Images



