

SUN2 Rabbit mAb

Catalog # AP78075

Specification

SUN2 Rabbit mAb - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW

WB, IHC-P, FC, ICC <u>Q9UH99</u> Human, Mouse, Rat Rabbit Monoclonal Antibody 80311

SUN2 Rabbit mAb - Additional Information

Gene ID 25777

Other Names SUN2

DilutionWB~~1/500-1/1000
IHC-P~~N/A
FC~~1:10~50
ICC~~N/A

Format

10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.

Storage

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

SUN2 Rabbit mAb - Protein Information

Name SUN2 (HGNC:14210)

Function

As a component of the LINC (LInker of Nucleoskeleton and Cytoskeleton) complex, involved in the connection between the nuclear lamina and the cytoskeleton. The nucleocytoplasmic interactions established by the LINC complex play an important role in the transmission of mechanical forces across the nuclear envelope and in nuclear movement and positioning. Specifically, SYNE2 and SUN2 assemble in arrays of transmembrane actin-associated nuclear (TAN) lines which are bound to F-actin cables and couple the nucleus to retrograde actin flow during actin-dependent nuclear movement. Required for interkinetic nuclear migration (INM) and essential for nucleokinesis and centrosome- nucleus coupling during radial neuronal migration in the cerebral cortex and during glial migration. Required for nuclear migration in retinal photoreceptor progenitors implicating association with cytoplasmic dynein-dynactin and kinesin motor complexes, and probably B-type lamins; SUN1 and SUN2 seem to act redundantly. The SUN1/2:KASH5 LINC complex couples telomeres to microtubules during meiosis; SUN1 and SUN2 seem to act at least partial





Tel: 858.875.1900 Fax: 858.875.1999

redundantly. Anchors chromosome movement in the prophase of meiosis and is involved in selective gene expression of coding and non-coding RNAs needed for gametogenesis. Required for telomere attachment to nuclear envelope and gametogenesis. May also function on endocytic vesicles as a receptor for RAB5-GDP and participate in the activation of RAB5.

Cellular Location

Nucleus inner membrane; Single-pass type II membrane protein. Nucleus envelope. Endosome membrane; Single-pass type II membrane protein

Tissue Location

Widely expressed. Highly expressed in heart, lung and muscle. Weakly expressed in fetal heart. Slightly overexpressed in some heart tissues form patients with congenital heart defects

SUN2 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

SUN2 Rabbit mAb - Images

