

GAPDH Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7873a

Specification

GAPDH Antibody (N-term) - Product Information

Application	IF, WB, IHC-P,E
Primary Accession	P04406
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	62-91

GAPDH Antibody (N-term) - Additional Information

Gene ID 2597

Other Names

Glyceraldehyde-3-phosphate dehydrogenase, GAPDH, Peptidyl-cysteine S-nitrosylase GAPDH, 2699-, GAPDH, GAPD

Target/Specificity

This GAPDH antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 62-91 amino acids from the N-terminal region of human GAPDH.

Dilution

IF~~1:10~50
WB~~1:1000
IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

GAPDH Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

GAPDH Antibody (N-term) - Protein Information

Name GAPDH {ECO:0000303|PubMed:2987855, ECO:0000312|HGNC:HGNC:4141}

Function Has both glyceraldehyde-3-phosphate dehydrogenase and nitrosylase activities, thereby

playing a role in glycolysis and nuclear functions, respectively (PubMed:[11724794](#), PubMed:[3170585](#)). Glyceraldehyde-3-phosphate dehydrogenase is a key enzyme in glycolysis that catalyzes the first step of the pathway by converting D- glyceraldehyde 3-phosphate (G3P) into 3-phospho-D-glyceroyl phosphate (PubMed:[11724794](#), PubMed:[3170585](#)). Modulates the organization and assembly of the cytoskeleton (By similarity). Facilitates the CHP1- dependent microtubule and membrane associations through its ability to stimulate the binding of CHP1 to microtubules (By similarity). Component of the GAIT (gamma interferon-activated inhibitor of translation) complex which mediates interferon-gamma-induced transcript-selective translation inhibition in inflammation processes (PubMed:[23071094](#)). Upon interferon-gamma treatment assembles into the GAIT complex which binds to stem loop-containing GAIT elements in the 3'-UTR of diverse inflammatory mRNAs (such as ceruplasmin) and suppresses their translation (PubMed:[23071094](#)). Also plays a role in innate immunity by promoting TNF-induced NF-kappa-B activation and type I interferon production, via interaction with TRAF2 and TRAF3, respectively (PubMed:[23332158](#), PubMed:[27387501](#)). Participates in nuclear events including transcription, RNA transport, DNA replication and apoptosis (By similarity). Nuclear functions are probably due to the nitrosylase activity that mediates cysteine S-nitrosylation of nuclear target proteins such as SIRT1, HDAC2 and PRKDC (By similarity).

Cellular Location


Cytoplasm, cytosol. Nucleus {ECO:0000250|UniProtKB:P04797}. Cytoplasm, perinuclear region. Membrane Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:P04797} Note=Translocates to the nucleus following S-nitrosylation and interaction with SIAH1, which contains a nuclear localization signal (By similarity). Postnuclear and Perinuclear regions (PubMed:12829261) {ECO:0000250|UniProtKB:P04797, ECO:0000269|PubMed:12829261}

GAPDH Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GAPDH Antibody (N-term) - Images




GAPDH Antibody (N-term) (Cat. # AP7873a) confocal immunofluorescent analysis with Hela cell.


0.025 mg/ml primary antibody was followed by FITC-conjugated goat anti-rabbit IgG (whole molecule). FITC emits green fluorescence. DAPI was used to stain the cell nuclear (blue).




Confocal immunofluorescent analysis of GAPDH Antibody (N-term)(Cat#AP7873a) with HeLa cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). Actin filaments have been labeled with Alexa Fluor 555 phalloidin (red). DAPI was used to stain the cell nuclear (blue).




All lanes : Anti-GAPDH Antibody (N-term) at 1:1000 dilution Lane 1: A431 whole cell lysate Lane 2: HeLa whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 36 kDa Blocking/Dilution buffer: 5% NFDN/TBST.



Western blot analysis of lysate from A375 cell line, using GAPDH Antibody (N-term)(Cat. #AP7873a). AP7873a was diluted at 1:500. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20ug.



Western blot analysis of lysates from HeLa, HUVEC cell line (from left to right), using GAPDH Antibody (N-term)(Cat. #AP7873a). AP7873a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.



Formalin-fixed and paraffin-embedded human hepatocarcinoma tissue reacted with GAPDH

antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

GAPDH Antibody (N-term) - Background

GAPDH catalyzes an important energy-yielding step in carbohydrate metabolism, the reversible oxidative phosphorylation of glyceraldehyde-3-phosphate in the presence of inorganic phosphate and nicotinamide adenine dinucleotide (NAD). The enzyme exists as a tetramer of identical chains.

GAPDH Antibody (N-term) - References

Azam,S., J. Biol. Chem. 283 (45), 30632-30641 (2008)
Lu,J., Biosci. Biotechnol. Biochem. 72 (9), 2432-2435 (2008)
Zhou,Y., Mol. Cancer Res. 6 (8), 1375-1384 (2008)

GAPDH Antibody (N-term) - Citations

- [Intermittent Bolus Compared With Continuous Feeding Enhances Insulin and Amino Acid Signaling to Translation Initiation in Skeletal Muscle of Neonatal Pigs](#)
- [Downregulated ZNF132 predicts unfavorable outcomes in breast Cancer via Hypermethylation modification](#)
- [The role of miR-128-3p through MAPK14 activation in the apoptosis of GC2 spermatocyte cell line following heat stress](#)
- [SHOC2 is associated with the survival of breast cancer cells and has prognostic value for patients with breast cancer](#)
- [MiR-29b/Sp1/FUT4 axis modulates the malignancy of leukemia stem cells by regulating fucosylation via Wnt/ \$\beta\$ -catenin pathway in acute myeloid leukemia.](#)
- [LINC01296/miR-26a/GALNT3 axis contributes to colorectal cancer progression by regulating O-glycosylated MUC1 via PI3K/AKT pathway.](#)
- [Wnt pathway is involved in 5-FU drug resistance of colorectal cancer cells.](#)
- [Atorvastatin ameliorates early brain injury through inhibition of apoptosis and ER stress in a rat model of subarachnoid hemorrhage.](#)
- [PSMD7 downregulation induces apoptosis and suppresses tumorigenesis of esophageal squamous cell carcinoma the mTOR/p70S6K pathway.](#)
- [Migration ability and Toll-like receptor expression of human mesenchymal stem cells improves significantly after three-dimensional culture.](#)
- [Proteasome inhibitor MG132 induces thyroid cancer cell apoptosis by modulating the activity of transcription factor FOXO3a.](#)
- [microRNA -140-5p inhibits colorectal cancer invasion and metastasis by targeting ADAMTS5 and IGFBP5.](#)
- [Increased expression of EHF via gene amplification contributes to the activation of HER family signaling and associates with poor survival in gastric cancer.](#)
- [Transient scrotal hyperthermia affects human sperm DNA integrity, sperm apoptosis, and sperm protein expression.](#)
- [The Ring Finger Protein RNF6 Induces Leukemia Cell Proliferation as a Direct Target of Pre-B-cell Leukemia Homeobox 1.](#)
- [miR-221/222 enhance the tumorigenicity of human breast cancer stem cells via modulation of PTEN/Akt pathway.](#)
- [The effect of 3-bromopyruvate on human colorectal cancer cells is dependent on glucose concentration but not hexokinase II expression.](#)
- [Prognostic significance of FAM3C in esophageal squamous cell carcinoma.](#)
- [Angiopoietin-like 4 enhances metastasis and inhibits apoptosis via inducing bone morphogenetic protein 7 in colorectal cancer cells.](#)
- [Elevated kinesin family member 26B is a prognostic biomarker and a potential therapeutic target for colorectal cancer.](#)
- [Oxidized low-density lipoprotein is associated with advanced-stage prostate cancer.](#)
- [Cell killing and radiosensitizing effects of atorvastatin in PC3 prostate cancer cells.](#)

- [MicroRNAs are involved in erythroid differentiation control.](#)