

**HuR Rabbit mAb**  
Catalog # AP79009**Specification****HuR Rabbit mAb - Product Information**

Application	WB, IHC-P, FC, IP, ICC
Primary Accession	<a href="#">Q15717</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	36092

**HuR Rabbit mAb - Additional Information****Gene ID** 1994**Other Names**  
ELAVL1**Dilution**  
WB~~1/500-1/1000  
IHC-P~~N/A  
FC~~1:10~50  
IP~~N/A  
ICC~~N/A**Format**  
10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.**Storage**  
Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.**HuR Rabbit mAb - Protein Information****Name** ELAVL1**Synonyms** HUR**Function**

RNA-binding protein that binds to the 3'-UTR region of mRNAs and increases their stability (PubMed: [14517288](http://www.uniprot.org/citations/14517288), PubMed: [18285462](http://www.uniprot.org/citations/18285462), PubMed: [31358969](http://www.uniprot.org/citations/31358969)). Involved in embryonic stem cell (ESC) differentiation: preferentially binds mRNAs that are not methylated by N6-methyladenosine (m6A), stabilizing them, promoting ESC differentiation (By similarity). Has also been shown to be capable of binding to m6A-containing mRNAs and contributes to MYC stability by binding to m6A-containing MYC mRNAs (PubMed: [32245947](http://www.uniprot.org/citations/32245947)). Binds to

poly-U elements and AU-rich elements (AREs) in the 3'-UTR of target mRNAs (PubMed:<a href="http://www.uniprot.org/citations/14731398" target="\_blank">14731398</a>, PubMed:<a href="http://www.uniprot.org/citations/17632515" target="\_blank">17632515</a>, PubMed:<a href="http://www.uniprot.org/citations/18285462" target="\_blank">18285462</a>, PubMed:<a href="http://www.uniprot.org/citations/23519412" target="\_blank">23519412</a>, PubMed:<a href="http://www.uniprot.org/citations/8626503" target="\_blank">8626503</a>). Binds avidly to the AU-rich element in FOS and IL3/interleukin-3 mRNAs. In the case of the FOS AU-rich element, binds to a core element of 27 nucleotides that contain AUUUA, AUUUUA, and AUUUUUA motifs. Binds preferentially to the 5'-UUUU[AG]UUU-3' motif in vitro (PubMed:<a href="http://www.uniprot.org/citations/8626503" target="\_blank">8626503</a>). With ZNF385A, binds the 3'-UTR of p53/TP53 mRNA to control their nuclear export induced by CDKN2A. Hence, may regulate p53/TP53 expression and mediate in part the CDKN2A anti-proliferative activity. May also bind with ZNF385A the CCNB1 mRNA (By similarity). Increases the stability of the leptin mRNA harboring an AU-rich element (ARE) in its 3' UTR (PubMed:<a href="http://www.uniprot.org/citations/29180010" target="\_blank">29180010</a>).

### Cellular Location

Cytoplasm. Nucleus. Cytoplasm, Stress granule {ECO:0000250|UniProtKB:P70372}. Cytoplasm, P-body. Note=Translocates into the cytoplasm following phosphorylation by MAPKAPK2 (PubMed:14517288). Likewise, phosphorylation by PRKCD promotes translocation from the nucleus into the cytoplasm, where it is associated with free and cytoskeleton-bound polysomes (PubMed:18285462). Localizes to the stress granules in the presence of PLEKHN1 (By similarity). {ECO:0000250|UniProtKB:P70372, ECO:0000269|PubMed:14517288, ECO:0000269|PubMed:18285462}

### Tissue Location

Ubiquitous. Detected in brain, liver, thymus and muscle.

### HuR Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### HuR Rabbit mAb - Images



