

**MEK1 (MAP2K1) Antibody (S218)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP7960D****Specification**

---

**MEK1 (MAP2K1) Antibody (S218) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q02750</a>
Other Accession	<a href="#">P36506</a> , <a href="#">Q63932</a> , <a href="#">P36507</a> , <a href="#">Q90891</a> , <a href="#">Q05116</a> , <a href="#">Q01986</a> , <a href="#">P29678</a> , <a href="#">P31938</a> , <a href="#">Q63980</a> , <a href="#">Q24324</a>
Reactivity	Human
Predicted	Drosophila, Hamster, Mouse, Rabbit, Rat, Xenopus, Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	43439
Antigen Region	196-225

**MEK1 (MAP2K1) Antibody (S218) - Additional Information****Gene ID** 5604**Other Names**

Dual specificity mitogen-activated protein kinase kinase 1, MAP kinase kinase 1, MAPKK 1, MKK1, ERK activator kinase 1, MAPK/ERK kinase 1, MEK 1, MAP2K1, MEK1, PRKMK1

**Target/Specificity**

This MEK1(MAP2K1) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 196-225 amino acids from human MEK1(MAP2K1).

**Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

MEK1 (MAP2K1) Antibody (S218) is for research use only and not for use in diagnostic or therapeutic procedures.

**MEK1 (MAP2K1) Antibody (S218) - Protein Information**

**Name** MAP2K1 ([HGNC:6840](#))

**Synonyms** MEK1, PRKMK1

**Function** Dual specificity protein kinase which acts as an essential component of the MAP kinase signal transduction pathway. Binding of extracellular ligands such as growth factors, cytokines and hormones to their cell-surface receptors activates RAS and this initiates RAF1 activation. RAF1 then further activates the dual-specificity protein kinases MAP2K1/MEK1 and MAP2K2/MEK2. Both MAP2K1/MEK1 and MAP2K2/MEK2 function specifically in the MAPK/ERK cascade, and catalyze the concomitant phosphorylation of a threonine and a tyrosine residue in a Thr-Glu-Tyr sequence located in the extracellular signal-regulated kinases MAPK3/ERK1 and MAPK1/ERK2, leading to their activation and further transduction of the signal within the MAPK/ERK cascade. Activates BRAF in a KSR1 or KSR2-dependent manner; by binding to KSR1 or KSR2 releases the inhibitory intramolecular interaction between KSR1 or KSR2 protein kinase and N-terminal domains which promotes KSR1 or KSR2-BRAF dimerization and BRAF activation (PubMed:[29433126](#)). Depending on the cellular context, this pathway mediates diverse biological functions such as cell growth, adhesion, survival and differentiation, predominantly through the regulation of transcription, metabolism and cytoskeletal rearrangements. One target of the MAPK/ERK cascade is peroxisome proliferator-activated receptor gamma (PPARG), a nuclear receptor that promotes differentiation and apoptosis. MAP2K1/MEK1 has been shown to export PPARG from the nucleus. The MAPK/ERK cascade is also involved in the regulation of endosomal dynamics, including lysosome processing and endosome cycling through the perinuclear recycling compartment (PNRC), as well as in the fragmentation of the Golgi apparatus during mitosis.

#### **Cellular Location**

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, microtubule organizing center, spindle pole body. Cytoplasm. Nucleus Membrane; Peripheral membrane protein. Note=Localizes at centrosomes during prometaphase, midzone during anaphase and midbody during telophase/cytokinesis (PubMed:14737111). Membrane localization is probably regulated by its interaction with KSR1 (PubMed:10409742)

#### **Tissue Location**


Widely expressed, with extremely low levels in brain.


### **MEK1 (MAP2K1) Antibody (S218) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **MEK1 (MAP2K1) Antibody (S218) - Images**





Western blot analysis of MAP2K1 (arrow) using rabbit polyclonal MAP2K1 Antibody (S217) (Cat.#AP7960d).293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the MAP2K1 gene (Lane 2) (Origene Technologies).



All lanes : Anti-MAP2K1 Antibody at 1:1000 dilution Lane 1: HepG2 whole cell lysate Lane 2: MCF-7 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 43 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

### MEK1 (MAP2K1) Antibody (S218) - Background

MAP2K1 is a member of the dual specificity protein kinase family, which acts as a mitogen-activated protein (MAP) kinase kinase. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This protein kinase lies upstream of MAP kinases and stimulates the enzymatic activity of MAP kinases upon wide variety of extra- and intracellular signals. As an essential component of MAP kinase signal transduction pathway, this kinase is involved in many cellular processes such as proliferation, differentiation, transcription regulation and development.

### MEK1 (MAP2K1) Antibody (S218) - References

Zheng, B., et al., Blood 102(3):1019-1027 (2003).

Li, S.P., et al., Cancer Res. 63(13):3473-3477 (2003).  
Zhu, X., et al., J. Neurochem. 86(1):136-142 (2003).  
Fringer, J., et al., J. Biol. Chem. 278(23):20612-20617 (2003).  
Witowsky, J.A., et al., J. Biol. Chem. 278(3):1403-1406 (2003).