

SEMA3E Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7976b**Specification**

SEMA3E Antibody (C-term) - Product Information

| | |
|-------------------|------------------------|
| Application | WB, IHC-P,E |
| Primary Accession | P70275 |
| Reactivity | Human, Mouse |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Antigen Region | 737-766 |

SEMA3E Antibody (C-term) - Additional Information**Gene ID** 20349**Other Names**

Semaphorin-3E, Semaphorin-H, Sema H, Sema3e, Semah, Semh

Target/Specificity

This SEMA3E antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 737-766 amino acids from the C-terminal region of human SEMA3E.

Dilution

WB~~1:1000

IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SEMA3E Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

SEMA3E Antibody (C-term) - Protein Information**Name** Sema3e**Synonyms** Semah, Semh**Function** Plays an important role in signaling via the cell surface receptor PLXND1. Mediates

reorganization of the actin cytoskeleton, leading to the retraction of cell projections. Promotes focal adhesion disassembly and inhibits adhesion of endothelial cells to the extracellular matrix. Regulates angiogenesis, both during embryogenesis and after birth. Can down-regulate sprouting angiogenesis. Required for normal vascular patterning during embryogenesis. Plays an important role in ensuring the specificity of synapse formation.

Cellular Location

Secreted.

Tissue Location


Detected in neurons in the thalamus. Detected in embryonic vasculature. Developing lungs, developing skeletal elements and ventral horns of the developing neural tube. Correlates positively with tumor progression.

SEMA3E Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


SEMA3E Antibody (C-term) - Images



Western blot analysis of Semaphorin 3E(Sema3e) (C-term) (Cat. #AP7976b) in CEM cell line lysates (35ug/lane).Sema3e (arrow) was detected using the purified Pab.



Western blot analysis of Semaphorin 3E(Sema3e) (C-term) (Cat. #AP7976b) in mouse lung tissue lysates (35ug/lane). Sema3e (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human breast carcinoma tissue reacted with Semaphorin 3E(Sema3e) (C-term) (Cat.#AP7976b), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

SEMA3E Antibody (C-term) - Background

The Semaphorins make up the largest family of axon guidance cues yet described. Semaphorins are divided into 8 classes (classes 3-7 found in vertebrates). Class 3 Semaphorins are secreted, classes 4 through 6 are transmembrane proteins, and class 7 are membrane associated via glycosylphosphatidylinositol (GPI) linkage. They are characterized structurally by a conserved ~400 amino acid sema domain. They are classically described as collapsing factors and mediators of axon repulsion, although they may also act as context-dependent chemoattractants. Semaphorins have been shown to have roles in cardiovascular development and in the regulation of immune cell antigen presentation. Receptors or receptor complexes that mediate semaphorin signaling include proteins of the Neuropilin and Plexin families.

SEMA3E Antibody (C-term) - References

Christensen C.R.L.,Cancer Res. 58:1238-1244(1998).

SEMA3E Antibody (C-term) - Citations

- [Cross-species Transcriptome Profiling Identifies New Alveolar Epithelial Type I Cell-specific Genes.](#)