

Endophilin B1 Antibody (Y80)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8034a

Specification

Endophilin B1 Antibody (Y80) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q9Y371
Other Accession	Q6AYE2 , Q9JK48 , Q32LM0
Reactivity	Human
Predicted	Bovine, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	40796
Antigen Region	59-86

Endophilin B1 Antibody (Y80) - Additional Information

Gene ID 51100

Other Names

Endophilin-B1, Bax-interacting factor 1, Bif-1, SH3 domain-containing GRB2-like protein B1, SH3GLB1, KIAA0491

Target/Specificity

This Endophilin B1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 59-86 amino acids from human Endophilin B1.

Dilution

WB~~1:1000

IHC-P~~1:10~50

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Endophilin B1 Antibody (Y80) is for research use only and not for use in diagnostic or therapeutic procedures.

Endophilin B1 Antibody (Y80) - Protein Information

Name SH3GLB1

Synonyms KIAA0491

Function May be required for normal outer mitochondrial membrane dynamics (PubMed:[15452144](#)). Required for coatamer-mediated retrograde transport in certain cells (By similarity). May recruit other proteins to membranes with high curvature. May promote membrane fusion (PubMed:[11604418](#)). Involved in activation of caspase-dependent apoptosis by promoting BAX/BAK1 activation (PubMed:[16227588](#)). Isoform 1 acts proapoptotic in fibroblasts (By similarity). Involved in caspase- independent apoptosis during nutrition starvation and involved in the regulation of autophagy. Activates lipid kinase activity of PIK3C3 during autophagy probably by associating with the PI3K complex II (PI3KC3-C2) (PubMed:[17891140](#)). Associated with PI3KC3-C2 during autophagy may regulate the trafficking of ATG9A from the Golgi complex to the peripheral cytoplasm for the formation of autophagosomes by inducing Golgi membrane tubulation and fragmentation (PubMed:[21068542](#)). Involved in regulation of degradative endocytic trafficking and cytokinesis, probably in the context of PI3KC3-C2 (PubMed:[20643123](#)). Isoform 2 acts antiapoptotic in neuronal cells; involved in maintenance of mitochondrial morphology and promotes neuronal viability (By similarity).

Cellular Location

Cytoplasm. Golgi apparatus membrane; Peripheral membrane protein. Mitochondrion outer membrane; Peripheral membrane protein. Cytoplasmic vesicle, autophagosome membrane. Midbody. Note=Association with the Golgi apparatus depends on the cell type (By similarity). Following starvation colocalizes with ATG5 and LC3 autophagy-related protein(s) on autophagosomal membranes (PubMed:17891140). {ECO:0000250, ECO:0000269|PubMed:17891140}

Tissue Location


Highly expressed in heart, skeletal muscle, kidney and placenta. Detected at lower levels in brain, colon, thymus, spleen, liver, small intestine, lung and peripheral blood leukocytes


Endophilin B1 Antibody (Y80) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Endophilin B1 Antibody (Y80) - Images






Western blot analysis of anti-Endophilin B1 Antibody (Y80) (Cat.#AP8034a) in HepG2 cell line lysates (35ug/lane). SH3GLB1 (arrow) was detected using the purified Pab.



Anti-Endophilin Antibody (Y80) at 1:1000 dilution + U118MG whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 41 kDa Blocking/Dilution buffer: 5% NFD/MTBST.



Formalin-fixed and paraffin-embedded human hepatocarcinoma tissue reacted with Endophilin B1 Antibody (Y80) (Cat.#AP8034a), which was peroxidase-conjugated to the secondary antibody,

followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

Endophilin B1 Antibody (Y80) - Background

Endophilin B1 may be required for normal outer mitochondrial membrane dynamics. It is required for coatamer-mediated retrograde transport in certain cells. It may recruit other proteins to membranes with high curvature and may promote membrane fusion.

Endophilin B1 Antibody (Y80) - References

Yamaguchi,H., J. Biol. Chem. 283 (27), 19112-19118 (2008)

Lee,J.W., Pathology 38 (4), 312-315 (2006)

Masuda,M., EMBO J. 25 (12), 2889-2897 (2006)