

MLKLAK Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8068c**Specification**

MLKLAK Antibody (Center) - Product Information

Application	IHC-P, WB,E
Primary Accession	O9NYL2
Other Accession	O9ESL4 , NP_598407
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	91155
Antigen Region	271-300

MLKLAK Antibody (Center) - Additional Information**Gene ID** 51776**Other Names**

Mitogen-activated protein kinase kinase kinase MLT, Human cervical cancer suppressor gene 4 protein, HCCS-4, Leucine zipper- and sterile alpha motif-containing kinase, MLK-like mitogen-activated protein triple kinase, Mixed lineage kinase-related kinase, MLK-related kinase, MRK, Sterile alpha motif- and leucine zipper-containing kinase AZK, ZAK, MLTK

Target/Specificity

This MLKLAK antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 271-300 amino acids from the Central region of human MLKLAK.

Dilution

IHC-P~~1:50~100

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MLKLAK Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

MLKLAK Antibody (Center) - Protein Information

Name MAP3K20 ([HGNC:17797](#))

Function Stress-activated component of a protein kinase signal transduction cascade that promotes programmed cell death in response to various stress, such as ribosomal stress, osmotic shock and ionizing radiation (PubMed:[10924358](#), PubMed:[11836244](#), PubMed:[12220515](#), PubMed:[14521931](#), PubMed:[15350844](#), PubMed:[15737997](#), PubMed:[18331592](#), PubMed:[20559024](#), PubMed:[26999302](#), PubMed:[32289254](#), PubMed:[32610081](#), PubMed:[35857590](#)). Acts by catalyzing phosphorylation of MAP kinase kinases, leading to activation of the JNK (MAPK8/JNK1, MAPK9/JNK2 and/or MAPK10/JNK3) and MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14) pathways (PubMed:[11042189](#), PubMed:[11836244](#), PubMed:[12220515](#), PubMed:[14521931](#), PubMed:[15172994](#), PubMed:[15737997](#), PubMed:[32289254](#), PubMed:[32610081](#), PubMed:[35857590](#)). Activates JNK through phosphorylation of MAP2K4/MKK4 and MAP2K7/MKK7, and MAP kinase p38 gamma (MAPK12) via phosphorylation of MAP2K3/MKK3 and MAP2K6/MKK6 (PubMed:[11836244](#), PubMed:[12220515](#)). Involved in stress associated with adrenergic stimulation: contributes to cardiac decompensation during periods of acute cardiac stress (PubMed:[15350844](#), PubMed:[21224381](#), PubMed:[27859413](#)). May be involved in regulation of S and G2 cell cycle checkpoint by mediating phosphorylation of CHEK2 (PubMed:[15342622](#)).

Cellular Location

Cytoplasm. Nucleus. Note=Translocates to the nucleus upon ultraviolet B irradiation.

Tissue Location


Ubiquitously expressed. Isoform ZAKbeta is the predominant form in all tissues examined, except for liver, in which isoform ZAKalpha is more highly expressed


MLKLAK Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MLKLAK Antibody (Center) - Images





Western blot analysis of anti-MLKLAK-C284 Pab (Cat. #AP8068c) in HepG2 cell line lysate (35ug/lane). MLKLAK-C284 (arrow) was detected using the purified Pab. This western blot detected isoform 2 of MLKLAK, the accession number for MLKLAK isoform 1 is Q9NYL2, NP_057737.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

MLKLAK Antibody (Center) - Background

MLKLAK is a member of the MAPKKK family of signal transduction molecules. It possesses an N-terminal kinase catalytic domain, followed by a leucine zipper motif and a sterile-alpha motif (SAM). This magnesium-binding protein forms homodimers and is located in the cytoplasm. The protein mediates gamma radiation signaling leading to cell cycle arrest and activity of this protein plays a role in cell cycle checkpoint regulation in cells. The protein also has pro-apoptotic activity.

MLKLAK Antibody (Center) - References

Blume-Jensen P, et al. Nature 2001. 411: 355.
Cantrell D, J. Cell Sci. 2001. 114: 1439.
Jiang S Oncogene 2000. 19: 5590.
Manning G, et al. Science 2002. 298: 1912.
Moller, D, et al. Am. J. Physiol. 1994. 266: C351-C359.

Robertson, S. et al. Trends Genet. 2000. 16: 368.
Robinson D, et al. Oncogene 2000. 19: 5548.
Van der Ven, P, et al. Hum. Molec. Genet. 1993. 2: 1889.
Vanhaesebroeck, B, et al. Biochem. J. 2000. 346: 561.
Van Weering D, et al. Recent Results Cancer Res. 1998. 154: 271.

MLKLAK Antibody (Center) - Citations

- [NEMO regulates a cell death switch in TNF signaling by inhibiting recruitment of RIPK3 to the cell death-inducing complex II.](#)