

PBP Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP8092c

Specification

PBP Antibody (Center) - Product Information

Application WB, IHC-P,E Primary Accession P30086

Reactivity Human, Mouse

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 21057
Antigen Region 137-167

PBP Antibody (Center) - Additional Information

Gene ID 5037

Other Names

Phosphatidylethanolamine-binding protein 1, PEBP-1, HCNPpp, Neuropolypeptide h3, Prostatic-binding protein, Raf kinase inhibitor protein, RKIP, Hippocampal cholinergic neurostimulating peptide, HCNP, PEBP1, PBP, PEBP

Target/Specificity

This PBP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 137-167 amino acids from the Central region of human PBP.

Dilution

WB~~1:1000 IHC-P~~1:50~100

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PBP Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

PBP Antibody (Center) - Protein Information





Name PEBP1

Synonyms PBP, PEBP

Function Binds ATP, opioids and phosphatidylethanolamine. Has lower affinity for phosphatidylinositol and phosphatidylcholine. Serine protease inhibitor which inhibits thrombin, neuropsin and chymotrypsin but not trypsin, tissue type plasminogen activator and elastase (By similarity). Inhibits the kinase activity of RAF1 by inhibiting its activation and by dissociating the RAF1/MEK complex and acting as a competitive inhibitor of MEK phosphorylation.


Cellular Location Cytoplasm.

PBP Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

PBP Antibody (Center) - Images




Western blot analysis of anti-PBP Antibody (Center) Antibody (Center) (Cat.#AP8092c) in 293 cell line lysates (35ug/lane). PBP(arrow) was detected using the purified Pab.






Western blot analysis of anti-PBP Antibody (Center) (Cat.#AP8092c) in mouse NIH-3T3 tissue lysates (35ug/lane). PBP(arrow) was detected using the purified Pab.



Western blot analysis of PBP (arrow) using rabbit polyclonal PBP Antibody (A152) (Cat. #AP8092c). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the PBP gene.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been



evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

PBP Antibody (Center) - Background

PBP binds ATP, opioids and phosphatidylethanolamine, exhibiting a lower affinity for phosphatidylinositol and phosphatidylcholine. This serine protease inhibitor inhibits thrombin, neuropsin and chymotrypsin but not trypsin, tissue type plasminogen activator and elastase. PBP contains hippocampal cholinergic neurostimulating peptide (HCNP), which may be involved in the function of the presynaptic cholinergic neurons of the central nervous system. HCNP increases the production of choline acetyltransferase but not acetylcholinesterase.

PBP Antibody (Center) - References

Tohdoh, N., et al., Brain Res. Mol. Brain Res. 30(2):381-384 (1995). Hori, N., et al., Gene 140(2):293-294 (1994). Seddiqi, N., et al., J. Mol. Evol. 39(6):655-660 (1994). Moore, C., et al., Brain Res. Mol. Brain Res. 37 (1-2), 74-78 (1996).