

TLK1 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8101b**Specification**

TLK1 Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q9UKI8
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	720-750

TLK1 Antibody (C-term) - Additional Information**Gene ID** 9874**Other Names**

Serine/threonine-protein kinase tousled-like 1, PKU-beta, Tousled-like kinase 1, TLK1, KIAA0137

Target/Specificity

This TLK1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 720-750 amino acids from the C-terminal region of human TLK1.

Dilution

WB~~1:8000

IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TLK1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

TLK1 Antibody (C-term) - Protein Information**Name** TLK1**Synonyms** KIAA0137**Function** Rapidly and transiently inhibited by phosphorylation following the generation of DNA

double-stranded breaks during S-phase. This is cell cycle checkpoint and ATM-pathway dependent and appears to regulate processes involved in chromatin assembly. Isoform 3 phosphorylates and enhances the stability of the t-SNARE SNAP23, augmenting its assembly with syntaxin. Isoform 3 protects the cells from the ionizing radiation by facilitating the repair of DSBs. In vitro, phosphorylates histone H3 at 'Ser-10'.

Cellular Location

Nucleus


Tissue Location

Widely expressed. Present in fetal placenta, liver, kidney and pancreas but not heart or skeletal muscle. Also found in adult cell lines. Isoform 3 is ubiquitously expressed in all tissues examined.


TLK1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TLK1 Antibody (C-term) - Images


Western blot analysis of hTLK1-L687 (Cat. #AP8101b) in K562 cell line lysates (35ug/lane). TLK1 (arrow) was detected using the purified Pab.



All lanes : Anti-TLK1 Antibody (L687) at 1:1000 dilution Lane 1: A431 whole cell lysate Lane 2: HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 87 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-TLK1 Antibody (L687) at 1:8000 dilution Lane 1: A431 whole cell lysate Lane 2: HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 87 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody,

which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

TLK1 Antibody (C-term) - Background

The Tousled-like kinases, first described in Arabadopsis, are nuclear serine/threonine kinases that are potentially involved in the regulation of chromatin assembly.[supplied by OMIM]

TLK1 Antibody (C-term) - References

Krause, D.R., et al., Oncogene 22(38):5927-5937 (2003).
Cabaniols, J.P., et al., Mol. Biol. Cell 10(12):4033-4041 (1999).
Sillje, H.H., et al., EMBO J. 18(20):5691-5702 (1999).
Yamakawa, A., et al., Gene 202 (1-2), 193-201 (1997).

TLK1 Antibody (C-term) - Citations

- [Silencing of Tousled-like kinase 1 sensitizes cholangiocarcinoma cells to cisplatin-induced apoptosis.](#)