

**WNK1 Antibody (Center)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP8107c****Specification**

---

**WNK1 Antibody (Center) - Product Information**

Application	WB,E
Primary Accession	<a href="#">O9H4A3</a>
Other Accession	<a href="#">O9JIH7</a> , <a href="#">P83741</a> , <a href="#">Q4VBX9</a>
Reactivity	Human, Mouse
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	250794

**WNK1 Antibody (Center) - Additional Information****Gene ID** 65125**Other Names**

Serine/threonine-protein kinase WNK1, Erythrocyte 65 kDa protein, p65, Kinase deficient protein, Protein kinase lysine-deficient 1, Protein kinase with no lysine 1, hWNK1, WNK1, HSN2, KDP, KIAA0344, PRKWNK1

**Target/Specificity**

This WNK1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the center region of human WNK1.

**Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

WNK1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**WNK1 Antibody (Center) - Protein Information****Name** WNK1 {ECO:0000303|PubMed:11571656, ECO:0000312|HGNC:HGNC:14540}

**Function** Serine/threonine-protein kinase component of the WNK1- SPAK/OSR1 kinase cascade, which acts as a key regulator of blood pressure and regulatory volume increase by promoting ion influx (PubMed:[15883153](#), PubMed:[17190791](#), PubMed:[31656913](#), PubMed:[34289367](#), PubMed:[36318922](#)). WNK1 mediates regulatory volume increase in response to hyperosmotic stress by acting as a molecular crowding sensor, which senses cell shrinkage and mediates formation of a membraneless compartment by undergoing liquid-liquid phase separation (PubMed:[36318922](#)). The membraneless compartment concentrates WNK1 with its substrates, OXSR1/OSR1 and STK39/SPAK, promoting WNK1-dependent phosphorylation and activation of downstream kinases OXSR1/OSR1 and STK39/SPAK (PubMed:[15883153](#), PubMed:[16263722](#), PubMed:[17190791](#), PubMed:[19739668](#), PubMed:[21321328](#), PubMed:[22989884](#), PubMed:[25477473](#), PubMed:[34289367](#), PubMed:[36318922](#)). Following activation, OXSR1/OSR1 and STK39/SPAK catalyze phosphorylation of ion cotransporters SLC12A1/NKCC2, SLC12A2/NKCC1, SLC12A5/KCC2 and SLC12A6/KCC3, regulating their activity (PubMed:[16263722](#), PubMed:[21321328](#)). Phosphorylation of Na-K-Cl cotransporters SLC12A2/NKCC1 and SLC12A2/NKCC1 promote their activation and ion influx; simultaneously, phosphorylation of K-Cl cotransporters SLC12A5/KCC2 and SLC12A6/KCC3 inhibit their activity, blocking ion efflux (PubMed:[19665974](#), PubMed:[21321328](#)). Also acts as a regulator of angiogenesis in endothelial cells via activation of OXSR1/OSR1 and STK39/SPAK: activation of OXSR1/OSR1 regulates chemotaxis and invasion, while STK39/SPAK regulates endothelial cell proliferation (PubMed:[25362046](#)). Also acts independently of the WNK1- SPAK/OSR1 kinase cascade by catalyzing phosphorylation of other substrates, such as SYT2, PCF11 and NEDD4L (PubMed:[29196535](#)). Mediates phosphorylation of SYT2, regulating SYT2 association with phospholipids and membrane-binding (By similarity). Regulates mRNA export in the nucleus by mediating phosphorylation of PCF11, thereby decreasing the association between PCF11 and POLR2A/RNA polymerase II and promoting mRNA export to the cytoplasm (PubMed:[29196535](#)). Acts as a negative regulator of autophagy (PubMed:[27911840](#)). Required for the abscission step during mitosis, independently of the WNK1-SPAK/OSR1 kinase cascade (PubMed:[21220314](#)). May also play a role in actin cytoskeletal reorganization (PubMed:[10660600](#)). Also acts as a scaffold protein independently of its protein kinase activity: negatively regulates cell membrane localization of various transporters and channels, such as SLC4A4, SLC26A6, SLC26A9, TRPV4 and CFTR (By similarity). Involved in the regulation of epithelial Na(+) channel (ENaC) by promoting activation of SGK1 in a kinase-independent manner: probably acts as a scaffold protein that promotes the recruitment of SGK1 to the mTORC2 complex in response to chloride, leading to mTORC2-dependent phosphorylation and activation of SGK1 (PubMed:[36373794](#)). Acts as an assembly factor for the ER membrane protein complex independently of its protein kinase activity: associates with EMC2 in the cytoplasm via its amphipathic alpha-helix, and prevents EMC2 ubiquitination and subsequent degradation, thereby promoting EMC2 stabilization (PubMed:[33964204](#)).

#### Cellular Location

Cytoplasm. Nucleus. Cytoplasm, cytoskeleton, spindle. Note=Mediates formation and localizes to cytoplasmic membraneless compartment in response to hyperosmotic stress (PubMed:[36318922](#)). Also localizes to the nucleus (PubMed:[29196535](#)) Localizes to the mitotic spindle during mitosis (PubMed:[21220314](#))

#### Tissue Location

Widely expressed, with highest levels observed in the testis, heart, kidney and skeletal muscle [Isoform 3]: This isoform is kidney-specific and specifically expressed in the distal convoluted tubule (DCT) and connecting tubule (CNT) of the nephron.


#### WNK1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **WNK1 Antibody (Center) - Images**



Western blot analysis of anti-hWNK1-S158 Pab (Cat. #AP8107c) in, from left to right, T47D, A375, HeLa, and mouse kidney cell line lysate (35ug/lane). hWNK1-S158 (arrow) was detected using the purified Pab (1:60 dilution).

#### **WNK1 Antibody (Center) - Background**

The WNK1 gene encodes a cytoplasmic serine-threonine kinase expressed in distal nephron.[supplied by OMIM]

#### **WNK1 Antibody (Center) - References**

Xu, B.E., et al., J. Biol. Chem. 277(50):48456-48462 (2002).  
Verissimo, F., et al., Oncogene 20(39):5562-5569 (2001).  
Wilson, F.H., et al., Science 293(5532):1107-1112 (2001).  
Moore, T.M., et al., J. Biol. Chem. 275(6):4311-4322 (2000).