

Diacylglycerol kinase iota (DGKI) Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8123b

Specification

Diacylglycerol kinase iota (DGKI) Antibody (C-term) - Product Information

Application WB, IHC-P,E Primary Accession 075912

Reactivity Human, Mouse

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 116233
Antigen Region 890-920

Diacylglycerol kinase iota (DGKI) Antibody (C-term) - Additional Information

Gene ID 9162

Other Names

Diacylglycerol kinase iota, DAG kinase iota, Diglyceride kinase iota, DGK-iota, DGKI

Target/Specificity

This Diacylglycerol kinase iota (DGKI) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 890-920 amino acids from the C-terminal region of human Diacylglycerol kinase iota (DGKI).

Dilution

WB~~1:1000 IHC-P~~1:50~100

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Diacylglycerol kinase iota (DGKI) Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Diacylglycerol kinase iota (DGKI) Antibody (C-term) - Protein Information

Name DGKI (HGNC:2855)



Function Diacylglycerol kinase that converts diacylglycerol/DAG into phosphatidic acid/phosphatidate/PA and regulates the respective levels of these two bioactive lipids (PubMed:23949095, PubMed:9830018). Thereby, acts as a central switch between the signaling pathways activated by these second messengers with different cellular targets and opposite effects in numerous biological processes (Probable). Has probably no preference for any of the diacylglycerols in terms of the acyl chain composition, especially for the acyl chain at the sn-2 position (PubMed:9830018). By controlling the diacylglycerol/DAG- mediated activation of RASGRP3, negatively regulates the Rap1 signaling pathway. May play a role in presynaptic diacylglycerol/DAG signaling and control neurotransmitter release during metabotropic glutamate receptor-dependent long-term depression (By similarity).

Cellular Location

Cell projection, axon {ECO:0000250|UniProtKB:F1MAB7}. Cell projection, dendrite {ECO:0000250|UniProtKB:F1MAB7}. Presynapse {ECO:0000250|UniProtKB:F1MAB7}. Postsynapse {ECO:0000250|UniProtKB:F1MAB7}. Postsynaptic density {ECO:0000250|UniProtKB:F1MAB7}. Synaptic cell membrane {ECO:0000250|UniProtKB:F1MAB7}. Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane {ECO:0000250|UniProtKB:F1MAB7}. Cytoplasm, cytosol. Nucleus. Note=Excluded from inhibitory synapses (By similarity). Localization between cytoplasm and nucleus is regulated by protein kinase C (PubMed:9830018). Both in the detergent soluble and particulate fractions (By similarity) {ECO:0000250|UniProtKB:F1MAB7, ECO:0000269|PubMed:9830018}

Tissue Location

Specifically expressed in brain and retina (PubMed:9830018). In brain, highly expressed in hippocampus, caudate nucleus, occipital pole, cerebral cortex, and cerebellum (PubMed:9830018). Also detected in kidney (PubMed:15894621)


Diacylglycerol kinase iota (DGKI) Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Diac	lalvcerol	kinase	iota	(DGKI)	Antibody	/ ((C-term)) -	Imag	es






The anti-DGKI Pab (Cat. #AP8123b) is used in Western blot to detect DGKI in mouse cerebellum tissue lysate.




DGKI Antibody (Y905) (Cat. #AP8123b) western blot analysis in A2058 cell line lysates (35ug/lane). This demonstrates the DGKI antibody detected the DGKI protein (arrow).



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

Diacylglycerol kinase iota (DGKI) Antibody (C-term) - Background





Tel: 858.875.1900 Fax: 858.875.1999

DGKI a member of the type IV diacylglycerol kinase subfamily. Diacylglycerol kinases regulate the intracellular concentration of diacylglycerol through its phosphorylation, producing phosphatidic acid. The specific role of the enzyme encoded by this gene is undetermined, however, it may play a crucial role in the production of phosphatidic acid in the retina or in recessive forms of retinal degeneration.

Diacylglycerol kinase iota (DGKI) Antibody (C-term) - References

Ding, L., et al., J. Biol. Chem. 273(49):32746-32752 (1998). Bowne, S.J., et al., Mol. Vis. 6, 6-9 (2000).