

Fructose 6 Phosphate Kinase (PFKM) Antibody (C-term) Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP8137b

Specification

Fructose 6 Phosphate Kinase (PFKM) Antibody (C-term) - Product Information

Application Primary Accession Other Accession Reactivity Predicted Host Clonality Isotype Calculated MW Antigen Region IF, IHC-P, WB,E <u>P08237</u> <u>Q60HD9</u> Human Monkey Rabbit Polyclonal Rabbit IgG 85183 746-776

Fructose 6 Phosphate Kinase (PFKM) Antibody (C-term) - Additional Information

Gene ID 5213

Other Names

ATP-dependent 6-phosphofructokinase, muscle type {ECO:0000255|HAMAP-Rule:MF_03184}, ATP-PFK {ECO:0000255|HAMAP-Rule:MF_03184}, PFK-M, 27111 {ECO:0000255|HAMAP-Rule:MF_03184}, 6-phosphofructokinase type A, Phosphofructo-1-kinase isozyme A, PFK-A, Phosphohexokinase {ECO:0000255|HAMAP-Rule:MF_03184}, PFKM, PFKX

Target/Specificity

This Fructose 6 Phosphate Kinase (PFKM) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 746-776 amino acids from the C-terminal region of human Fructose 6 Phosphate Kinase (PFKM).

Dilution $IF \sim 1:10 \sim 50$ $IHC-P \sim 1:50 \sim 100$ $WB \sim 1:1000$ $E \sim Use$ at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Fructose 6 Phosphate Kinase (PFKM) Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.



Fructose 6 Phosphate Kinase (PFKM) Antibody (C-term) - Protein Information

Name PFKM

Synonyms PFKX

Function Catalyzes the phosphorylation of D-fructose 6-phosphate to fructose 1,6-bisphosphate by ATP, the first committing step of glycolysis.


Cellular Location Cytoplasm {ECO:0000255|HAMAP-Rule:MF_03184}.

Fructose 6 Phosphate Kinase (PFKM) Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Fructose 6 Phosphate Kinase (PFKM) Antibody (C-term) - Images



Confocal immunofluorescent analysis of Fructose 6 Phosphate Kinase (PFKM) Antibody(C-term)(Cat#AP8137b) with Hela cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).






Western blot analysis of PFKM polyclonal antibody (Cat. #AP8137b) in Hela cell line lysate. PFKM(arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human skeletal muscle reacted with PFKM Antibody (C-term)(Cat.#AP8137b), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Formalin-fixed and paraffin-embedded human hepatocarcinoma tissue reacted with PFKM antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

Fructose 6 Phosphate Kinase (PFKM) Antibody (C-term) - Background



Phosphofructokinase catalyzes the irreversible conversion of fructose 6 phosphate to fructose 1,6 bisphosphate. Mammalian PFK is a complex isozyme consisting of 3 subunits: muscle (M), liver (L), and platelet (P). Only M type PFK isozyme is expressed in mature muscle, while erythrocytes contain both L and M subunits. Defects in PFKM are the cause of glycogen storage disease type 7 (GSD7), also known as Tarui disease.

Fructose 6 Phosphate Kinase (PFKM) Antibody (C-term) - References

Howard, T.D., et al., Genomics 34(1):122-127 (1996). Vasconcelos, O., et al., Proc. Natl. Acad. Sci. U.S.A. 92(22):10322-10326 (1995). Raben, N., et al., J. Biol. Chem. 268(7):4963-4967 (1993). Yamasaki, T., et al., Gene 104(2):277-282 (1991). Sharma, P.M., et al., J. Biol. Chem. 265(16):9006-9010 (1990).