

**PFKFB2 Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP8146b**

**Specification**

---

**PFKFB2 Antibody (C-term) - Product Information**

|                   |   |
|-------------------|---|
| Application       | WB,E  |
| Primary Accession | <a href="#">O60825</a>                          |
| Other Accession   | <a href="#">O9JJH5</a> , <a href="#">P70265</a> |
| Reactivity        | Human   |
| Predicted         | Mouse, Rat                                      |
| Host              | Rabbit  |
| Clonality         | Polyclonal                                      |
| Isotype           | Rabbit IgG                                      |
| Antigen Region    | 467-497   |

**PFKFB2 Antibody (C-term) - Additional Information**

**Gene ID** 5208

**Other Names**

6-phosphofructo-2-kinase/fructose-2, 6-bisphosphatase 2, 6PF-2-K/Fru-2, 6-P2ase 2, PFK/FBPase 2, 6PF-2-K/Fru-2, 6-P2ase heart-type isozyme, 6-phosphofructo-2-kinase, Fructose-2, 6-bisphosphatase, PFKFB2

**Target/Specificity**

This PFKFB2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 467-497 amino acids from the C-terminal region of human PFKFB2.

**Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

PFKFB2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**PFKFB2 Antibody (C-term) - Protein Information**

**Name** PFKFB2 ([HGNC:8873](#))

**Function** Synthesis and degradation of fructose 2,6-bisphosphate.


**Tissue Location**  
Heart.

### **PFKFB2 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **PFKFB2 Antibody (C-term) - Images**



The anti-PFKFB2 Pab (Cat. #AP8146b) is used in Western blot to detect PFKFB2 in Jurkat cell lysate.

### **PFKFB2 Antibody (C-term) - Background**

Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the  $\gamma$  phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains. The AGC kinase group consists of 63 kinases including the cyclic nucleotide-regulated protein kinase (PKA & PKG) family, the diacylglycerol-activated/phospholipid-dependent protein kinase C (PKC) family, the related to PKA and PKC (RAC/Akt) protein kinase family, the kinases that phosphorylate G protein-coupled receptors family (ARK), and the kinases that phosphorylate ribosomal protein S6 family (RSK).

**PFKFB2 Antibody (C-term) - References**

Soejima, H., et al., Genomics 74(1):115-120 (2001).

Heine-Suner, D., et al., Eur. J. Biochem. 254(1):103-110 (1998).

**PFKFB2 Antibody (C-term) - Citations**

- [Akt-dependent activation of the heart 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase \(PFKFB2\) isoenzyme by amino acids.](#)