

## **HNF1A Antibody (Center)**

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8717c

# **Specification**

## **HNF1A Antibody (Center) - Product Information**

Application IF, WB,E Primary Accession P20823

Other Accession P15257, P22361
Reactivity Human, Mouse

Predicted Rat
Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 67386
Antigen Region 177-205

# **HNF1A** Antibody (Center) - Additional Information

#### **Gene ID 6927**

### **Other Names**

Hepatocyte nuclear factor 1-alpha, HNF-1-alpha, HNF-1A, Liver-specific transcription factor LF-B1, LFB1, Transcription factor 1, TCF-1, HNF1A, TCF1

## Target/Specificity

This HNF1A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 177-205 amino acids from the Central region of human HNF1A.

#### **Dilution**

IF~~1:10~50 WB~~1:1000

E~~Use at an assay dependent concentration.

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

## **Precautions**

HNF1A Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

#### **HNF1A Antibody (Center) - Protein Information**





#### Name HNF1A

#### **Synonyms TCF1**

**Function** Transcriptional activator that regulates the tissue specific expression of multiple genes, especially in pancreatic islet cells and in liver (By similarity). Binds to the inverted palindrome 5'-GTTAATNATTAAC-3' (PubMed: 10966642, PubMed: 12453420). Activates the transcription of CYP1A2, CYP2E1 and CYP3A11 (By similarity).

## **Cellular Location**

 $Nucleus \ \{ ECO: 0000255 | PROSITE-ProRule: PRU00108, ECO: 0000269 | PubMed: 10966642, ECO: 0000269 | PubMed: 38018242 \}$ 


Tissue Location Liver.

## **HNF1A Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# **HNF1A Antibody (Center) - Images**



Confocal immunofluorescent analysis of HNF1A Antibody (Center)(Cat#AP8717c) with HepG2 cell followed by Alexa Fluor® 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).





Western blot analysis of HNF1A Antibody (Center) (Cat. #AP8717c) in mouse liver tissue lysates (35ug/lane). HNF1A (arrow) was detected using the purified Pab.

# HNF1A Antibody (Center) - Background

HNF1A is a transcription factor required for the expression of several liver-specific genes. This protein functions as a homodimer and binds to the inverted palindrome 5'-GTTAATNATTAAC-3'.

# **HNF1A Antibody (Center) - References**

Bach, I., et.al., Genomics 8 (1), 155-164 (1990)