

**GTF2H4 Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP8730C**

**Specification**

---

**GTF2H4 Antibody (Center) - Product Information**

Application	WB,E
Primary Accession	<a href="#">O92759</a>
Other Accession	<a href="#">O70422</a>
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	52186
Antigen Region	123-151

**GTF2H4 Antibody (Center) - Additional Information**

**Gene ID** 2968

**Other Names**

General transcription factor IIH subunit 4, Basic transcription factor 2 52 kDa subunit, BTF2 p52, General transcription factor IIH polypeptide 4, TFIIH basal transcription factor complex p52 subunit, GTF2H4

**Target/Specificity**

This GTF2H4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 123-151 amino acids from the Central region of human GTF2H4.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

GTF2H4 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**GTF2H4 Antibody (Center) - Protein Information**

**Name** GTF2H4

**Function** Component of the general transcription and DNA repair factor IIH (TFIIH) core complex, which is involved in general and transcription-coupled nucleotide excision repair (NER) of damaged DNA and, when complexed to CAK, in RNA transcription by RNA polymerase II. In NER, TFIIH acts by opening DNA around the lesion to allow the excision of the damaged oligonucleotide and its replacement by a new DNA fragment. In transcription, TFIIH has an essential role in transcription initiation. When the pre-initiation complex (PIC) has been established, TFIIH is required for promoter opening and promoter escape. Phosphorylation of the C-terminal tail (CTD) of the largest subunit of RNA polymerase II by the kinase module CAK controls the initiation of transcription.

#### **Cellular Location**


Nucleus.

### **GTF2H4 Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **GTF2H4 Antibody (Center) - Images**



Western blot analysis of GTF2H4 Antibody (Center) (Cat. #AP8730c) in Y79 cell line lysates (35ug/lane). GTF2H4 (arrow) was detected using the purified Pab.

### **GTF2H4 Antibody (Center) - Background**

Component of the core-TFIIH basal transcription factor involved in nucleotide excision repair (NER) of DNA and, when complexed to CAK, in RNA transcription by RNA polymerase II.

### **GTF2H4 Antibody (Center) - References**

Blau, J., et.al., Mol. Cell. Biol. 16 (5), 2044-2055 (1996)  
Zhou, Q. et.al., Science 274 (5287), 605-610 (1996)