

BDK_1 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8735C**Specification**

BDK_1 Antibody (Center) - Product Information

Application	WB, FC,E
Primary Accession	P46663
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	40495
Antigen Region	213-239

BDK_1 Antibody (Center) - Additional Information**Gene ID** 623**Other Names**

B1 bradykinin receptor, B1R, BK-1 receptor, BDKRB1, BRADYB1

Target/Specificity

This BDK_1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 213-239 amino acids from the Central region of human BDK_1.

Dilution

WB~~1:1000

FC~~1:10~50

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

BDK_1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

BDK_1 Antibody (Center) - Protein Information**Name** BDKRB1**Synonyms** BRADYB1

Function This is a receptor for bradykinin. Could be a factor in chronic pain and inflammation.

Cellular Location


Cell membrane; Multi-pass membrane protein

BDK_1 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BDK_1 Antibody (Center) - Images



All lanes : Anti-BDK_1 Antibody (Center) at 1:1000 dilution Lane 1: HepG2 whole cell lysate Lane 2: HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 40 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



BDKRB1 Antibody (Center) (Cat. #AP8735c) flow cytometric analysis of HepG2 cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

BDK_1 Antibody (Center) - Background

Bradykinin, a 9 aa peptide, is generated in pathophysiologic conditions such as inflammation, trauma, burns, shock, and allergy. Two types of G-protein coupled receptors have been found which bind bradykinin and mediate responses to these pathophysiologic conditions. BDKRB1 is one of these receptors and is synthesized de novo following tissue injury. Receptor binding leads to an increase in the cytosolic calcium ion concentration, ultimately resulting in chronic and acute inflammatory responses.

BDK_1 Antibody (Center) - References

Bachvarov, D.R., et.al., Genomics 33 (3), 374-381 (1996)