

HIST1H2AH Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8820a**Specification**

HIST1H2AH Antibody (N-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	Q96KK5
Other Accession	P02263 , Q4FZT6 , Q8BFU2 , Q7L7L0 , P35062 , Q64523 , Q16777 , A1A4R1 , Q64522 , Q8IUE6 , P0CC09 , Q6GSS7 , Q6FI13 , P02262 , P22752 , P0C0S8 , P0C0S9 , Q8CGP7 , Q99878 , Q8CGP6 , Q64598 , Q8CGP5 , P0C170 , P20671 , P0C169 , Q93077 , P04908 , NP_542163 , C0HKE1 , C0HKE2 , C0HKE3 , C0HK
Reactivity	Human
Predicted	Mouse, Rat, Bovine, Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	13906
Antigen Region	1-30

HIST1H2AH Antibody (N-term) - Additional Information**Gene ID** 85235**Other Names**

Histone H2A type 1-H, Histone H2A/s, HIST1H2AH, HIST1H2AI

Target/Specificity

This HIST1H2AH antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human HIST1H2AH.

Dilution

WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HIST1H2AH Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

HIST1H2AH Antibody (N-term) - Protein Information

Name H2AC12 ([HGNC:13671](#))

Function Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Cellular Location


Nucleus. Chromosome.

HIST1H2AH Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


HIST1H2AH Antibody (N-term) - Images



Western blot analysis of HIST1H2AH Antibody (N-term) (Cat. #AP8820a) in A2058, A375, K562, HeLa, MCF-7 cell line lysates (35ug/lane). HIST1H2AH (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human brain tissue reacted with HIST1H2AH Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



HIST1H2AH Antibody (N-term) (Cat. #AP8820a) flow cytometric analysis of MCF-7 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

HIST1H2AH Antibody (N-term) - Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a member of the histone H2A family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element.

HIST1H2AH Antibody (N-term) - References

Kimura, H., et al., J. Cell Biol. 175 (3), 389-400 (2006)