

**TCERG1L Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP8880c****Specification**

---

**TCERG1L Antibody (Center) - Product Information**

Application	FC, IHC-P, WB,E
Primary Accession	<a href="#">Q5VW11</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	65660
Antigen Region	392-421

**TCERG1L Antibody (Center) - Additional Information****Gene ID** 256536**Other Names**

Transcription elongation regulator 1-like protein, TCERG1L

**Target/Specificity**

This TCERG1L antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 392-421 amino acids from the Central region of human TCERG1L.

**Dilution**

FC~~1:10~50

IHC-P~~1:50~100

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

TCERG1L Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.


**TCERG1L Antibody (Center) - Protein Information****Name** TCERG1L

## TCERG1L Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


## TCERG1L Antibody (Center) - Images



Western blot analysis of TCERG1L Antibody (Center) (Cat. #AP8880c) in mouse stomach tissue lysates (35ug/lane). TCERG1L (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human brain tissue reacted with TCERG1L Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



TCERG1L Antibody (Center) (Cat. #AP8880c) flow cytometric analysis of 293 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

#### **TCERG1L Antibody (Center) - References**

Roeske, D., et al., Mol. Psychiatry (2009)