

KCNQ1 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8944c**Specification**

KCNQ1 Antibody (Center) - Product Information

Application	FC, WB,E
Primary Accession	P51787
Other Accession	Q9Z0N7 , P97414 , F1QG65 , Q9MYS6 , P70057 , Q9TTJ7
Reactivity	Human, Mouse
Predicted	Pig, Rabbit, Rat, Xenopus
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	74699
Antigen Region	513-540

KCNQ1 Antibody (Center) - Additional Information**Gene ID** 3784**Other Names**

Potassium voltage-gated channel subfamily KQT member 1, IKs producing slow voltage-gated potassium channel subunit alpha KvLQT1, KQT-like 1, Voltage-gated potassium channel subunit Kv71, KCNQ1, KCNA8, KCNA9, KVLQT1

Target/Specificity

This KCNQ1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 513-540 amino acids from the Central region of human KCNQ1.

Dilution

FC~~1:10~50

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

KCNQ1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

KCNQ1 Antibody (Center) - Protein Information

Name KCNQ1 ([HGNC:6294](#))

Function Pore-forming subunit of the voltage-gated potassium (Kv) channel involved in the regulation of cardiomyocyte excitability and important in normal development and functions of myocardium, inner ear, stomach and colon (PubMed:[10646604](#), PubMed:[25441029](#)). Associates with KCNE beta subunits that modulates current kinetics (PubMed:[10646604](#), PubMed:[11101505](#), PubMed:[19687231](#), PubMed:[8900283](#), PubMed:[9108097](#), PubMed:[9312006](#)). Induces a voltage-dependent current by rapidly activating and slowly deactivating potassium-selective outward current (PubMed:[10646604](#), PubMed:[11101505](#), PubMed:[25441029](#), PubMed:[8900283](#), PubMed:[9108097](#), PubMed:[9312006](#)). Also promotes a delayed voltage activated potassium current showing outward rectification characteristic (By similarity). During beta-adrenergic receptor stimulation, participates in cardiac repolarization by associating with KCNE1 to form the I(Ks) cardiac potassium current that increases the amplitude and slows down the activation kinetics of outward potassium current I(Ks) (By similarity) (PubMed:[10646604](#), PubMed:[11101505](#), PubMed:[8900283](#), PubMed:[9108097](#), PubMed:[9312006](#)). Muscarinic agonist oxotremorine-M strongly suppresses KCNQ1/KCNE1 current (PubMed:[10713961](#)). When associated with KCNE3, forms the potassium channel that is important for cyclic AMP-stimulated intestinal secretion of chloride ions (PubMed:[10646604](#)). This interaction with KCNE3 is reduced by 17beta-estradiol, resulting in the reduction of currents (By similarity). During conditions of increased substrate load, maintains the driving force for proximal tubular and intestinal sodium ions absorption, gastric acid secretion, and cAMP-induced jejunal chloride ions secretion (By similarity). Allows the provision of potassium ions to the luminal membrane of the secretory canaliculus in the resting state as well as during stimulated acid secretion (By similarity). When associated with KCNE2, forms a heterooligomer complex leading to currents with an apparently instantaneous activation, a rapid deactivation process and a linear current-voltage relationship and decreases the amplitude of the outward current (PubMed:[11101505](#)). When associated with KCNE4, inhibits voltage-gated potassium channel activity (PubMed:[19687231](#)). When associated with KCNE5, this complex only conducts current upon strong and continued depolarization (PubMed:[12324418](#)). Also forms a heterotetramer with KCNQ5; has a voltage-gated potassium channel activity (PubMed:[24855057](#)). Binds with phosphatidylinositol 4,5-bisphosphate (PubMed:[25037568](#)). KCNQ1-KCNE2 channel associates with Na(+)-coupled myo-inositol symporter in the apical membrane of choroid plexus epithelium and regulates the myo- inositol gradient between blood and cerebrospinal fluid with an impact on neuron excitability (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein. Cytoplasmic vesicle membrane Early endosome. Membrane raft. Endoplasmic reticulum Basolateral cell membrane. Apical cell membrane {ECO:0000250|UniProtKB:P97414}; Multi-pass membrane protein. Note=Colocalized with KCNE3 at the plasma membrane (PubMed:10646604). Upon 17beta-oestradiol treatment, colocalizes with RAB5A at early endosome (PubMed:23529131). Heterotetramer with KCNQ5 is highly retained at the endoplasmic reticulum and is localized outside of lipid raft microdomains (PubMed:24855057). During the early stages of epithelial cell polarization induced by the calcium switch, it is removed from the plasma membrane to the endoplasmic reticulum, where it is retained, and redistributed to the basolateral cell surface in a PI3K-dependent manner at a later stage (PubMed:21228319). Colocalizes with SLC5A3 at the apical membrane of choroid plexus epithelium {ECO:0000250|UniProtKB:P97414, ECO:0000269|PubMed:10646604, ECO:0000269|PubMed:21228319, ECO:0000269|PubMed:23529131, ECO:0000269|PubMed:24855057}

Tissue Location


Abundantly expressed in heart, pancreas, prostate, kidney, small intestine and peripheral blood leukocytes. Less abundant in placenta, lung, spleen, colon, thymus, testis and ovaries

KCNQ1 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KCNQ1 Antibody (Center) - Images



Western blot analysis of KCNQ1 Antibody (Center) (Cat. #AP8944c) in mouse lung tissue lysates (35ug/lane). KCNQ1 (arrow) was detected using the purified Pab.



KCNQ1 Antibody (Center) (Cat. #AP8944c) flow cytometric analysis of 293 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

KCNQ1 Antibody (Center) - Background

KCNQ1 is a protein for a voltage-gated potassium channel required for the repolarization phase of the cardiac action potential. The gene product can form heteromultimers with two other potassium channel proteins, KCNE1 and KCNE3.

KCNQ1 Antibody (Center) - References

Holm,H., et.al., Nat. Genet. 42 (2), 117-122 (2010)
Ohshige,T., et. al., Diabetes Care (2010) In press