

Phospho-PKA R2 (S99) Antibody

Rabbit mAb Catalog # AP90120

Specification

Phospho-PKA R2 (S99) Antibody - Product Information

Application WB, IHC, ICC, IP

Primary Accession
Reactivity
Rat, Pig
Clonality
Monoclonal

Other Names

KAP2; PRKAR2; PKR2;

Isotype Rabbit IgG
Host Rabbit
Calculated MW 45518 Da

Phospho-PKA R2 (S99) Antibody - Additional Information

Dilution WB~~1:1000

IHC~~1:100~500

ICC~~N/A IP~~N/A

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human

Phospho-PKA R2 (S99)

Description Regulatory subunit of the cAMP-dependent

protein kinases involved in cAMP signaling in cells. Type II regulatory chains mediate membrane association by binding to anchoring proteins, including the MAP2

kinase.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline ,

pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid

freeze / thaw cycle.

Phospho-PKA R2 (S99) Antibody - Protein Information

Name PRKAR2A

Synonyms PKR2, PRKAR2

Function

Regulatory subunit of the cAMP-dependent protein kinases involved in cAMP signaling in cells. Type II regulatory chains mediate membrane association by binding to anchoring proteins, including the MAP2 kinase.



Cellular Location

Cytoplasm. Cell membrane. Note=Colocalizes with PJA2 in the cytoplasm and the cell membrane

Tissue Location


Four types of regulatory chains are found: I-alpha, I-beta, II-alpha, and II-beta. Their expression varies among tissues and is in some cases constitutive and in others inducible

Phospho-PKA R2 (S99) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Phospho-PKA R2 (S99) Antibody - Images



Western blot analysis of Phospho-PKA R2 (Ser99) in K562 cell lysate.