

**Neuropilin 1 Antibody**  
**Rabbit mAb**  
**Catalog # AP90289****Specification**

---

**Neuropilin 1 Antibody - Product Information**

Application	WB, IHC, FC, ICC, IP
Primary Accession	<a href="#">O14786</a>
Reactivity	Rat
Clonality	Monoclonal

**Other Names**

A5 protein; BDCA4; CD304 ; Neuropilin1; NP1; NPN1; NRP1; VEGF165R ; NEL like protein 1; Blood dendritic cell antigen 4 ;

Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	103134 Da

**Neuropilin 1 Antibody - Additional Information**

Dilution	WB~~1:1000 IHC~~1:100~500 FC~~1:10~50 ICC~~N/A IP~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Neuropilin 1
Description	The membrane-bound isoform 1 is a receptor involved in the development of the cardiovascular system, in angiogenesis, in the formation of certain neuronal circuits and in organogenesis outside the nervous system. It mediates the chemorepulsant activity of semaphorins. It binds to semaphorin 3A, The PLGF-2 isoform of PGF, The VEGF-165 isoform of VEGF and VEGF-B. Coexpression with KDR results in increased VEGF-165 binding to KDR as well as increased chemotaxis. It may regulate VEGF-induced angiogenesis. The soluble isoform 2 binds VEGF-165 and appears to inhibit its binding to cells.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

## Neuropilin 1 Antibody - Protein Information

**Name** NRP1 ([HGNC:8004](#))

**Synonyms** NRP, VEGF165R

### Function

Cell-surface receptor involved in the development of the cardiovascular system, in angiogenesis, in the formation of certain neuronal circuits and in organogenesis outside the nervous system. Mediates the chemorepulsant activity of semaphorins (PubMed:<a href="http://www.uniprot.org/citations/10688880" target="\_blank">10688880</a>, PubMed:<a href="http://www.uniprot.org/citations/9288753" target="\_blank">9288753</a>, PubMed:<a href="http://www.uniprot.org/citations/9529250" target="\_blank">9529250</a>). Recognizes a C-end rule (CendR) motif R/KXXR/K on its ligands which causes cellular internalization and vascular leakage (PubMed:<a href="http://www.uniprot.org/citations/19805273" target="\_blank">19805273</a>). It binds to semaphorin 3A, the PLGF-2 isoform of PGF, the VEGF165 isoform of VEGFA and VEGFB (PubMed:<a href="http://www.uniprot.org/citations/10688880" target="\_blank">10688880</a>, PubMed:<a href="http://www.uniprot.org/citations/19805273" target="\_blank">19805273</a>, PubMed:<a href="http://www.uniprot.org/citations/9288753" target="\_blank">9288753</a>, PubMed:<a href="http://www.uniprot.org/citations/9529250" target="\_blank">9529250</a>). Coexpression with KDR results in increased VEGF165 binding to KDR as well as increased chemotaxis. Regulates VEGF-induced angiogenesis. Binding to VEGFA initiates a signaling pathway needed for motor neuron axon guidance and cell body migration, including for the caudal migration of facial motor neurons from rhombomere 4 to rhombomere 6 during embryonic development (By similarity). Regulates mitochondrial iron transport via interaction with ABCB8/MITOSUR (PubMed:<a href="http://www.uniprot.org/citations/30623799" target="\_blank">30623799</a>).

### Cellular Location

[Isoform 2]: Secreted

### Tissue Location


[Isoform 1]: The expression of isoforms 1 and 2 does not seem to overlap. Expressed in olfactory epithelium (at protein level) (PubMed:33082293). Expressed in fibroblasts (at protein level) (PubMed:36213313). Expressed by the blood vessels of different tissues In the developing embryo it is found predominantly in the nervous system. In adult tissues, it is highly expressed in heart and placenta; moderately in lung, liver, skeletal muscle, kidney and pancreas; and low in adult brain (PubMed:10688880, PubMed:9529250). Expressed in the central nervous system, including olfactory related regions such as the olfactory tubercles and paraolfactory gyri (PubMed:33082293)

## Neuropilin 1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Neuropilin 1 Antibody - Images



Western blot analysis of Neuropilin 1 expression in mouse heart lysate.