

**Phospho-p95/NBS1 (S343) Antibody**  
Rabbit mAb  
Catalog # AP90300**Specification****Phospho-p95/NBS1 (S343) Antibody - Product Information**

Application **WB, ICC, IP**  
Primary Accession **[O60934](#)**  
Clonality **Monoclonal**

**Other Names**

Cell cycle regulatory protein P95, NBN, NBS, NIBRIN, NIJMEGEN BREAKAGE syndrome protein 1, p95-NBS1

Isotype **Rabbit IgG**  
Host **Rabbit**  
Calculated MW **84959 Da**

**Phospho-p95/NBS1 (S343) Antibody - Additional Information**

Dilution **WB~~1:1000**  
**ICC~~N/A**  
**IP~~N/A**

Purification **Affinity-chromatography**  
Immunogen **A synthesized peptide derived from human Phospho-p95/NBS1 (S343)**

Description **NBS1 is a member of the MRE11/RAD50 double-strand break repair complex. Involved in DNA double-strand break repair and DNA damage-induced checkpoint activation. Mutation results in the Nijmegen breakage syndrome (NBS), an autosomal recessive chromosomal instability syndrome.**

Storage Condition and Buffer **Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.**

**Phospho-p95/NBS1 (S343) Antibody - Protein Information**

Name NBN ([HGNC:7652](#))

**Function**

Component of the MRN complex, which plays a central role in double-strand break (DSB) repair, DNA recombination, maintenance of telomere integrity and meiosis (PubMed:<a href="http://www.uniprot.org/citations/10888888" target="\_blank">10888888</a>, PubMed:<a href="http://www.uniprot.org/citations/15616588" target="\_blank">15616588</a>, PubMed:<a href="http://www.uniprot.org/citations/18411307" target="\_blank">18411307</a>, PubMed:<a

[18583988](http://www.uniprot.org/citations/18583988), PubMed:[18678890](http://www.uniprot.org/citations/18678890), PubMed:[19759395](http://www.uniprot.org/citations/19759395), PubMed:[23115235](http://www.uniprot.org/citations/23115235), PubMed:[28216226](http://www.uniprot.org/citations/28216226), PubMed:[28867292](http://www.uniprot.org/citations/28867292), PubMed:[9705271](http://www.uniprot.org/citations/9705271)). The MRN complex is involved in the repair of DNA double-strand breaks (DSBs) via homologous recombination (HR), an error-free mechanism which primarily occurs during S and G2 phases (PubMed:[19759395](http://www.uniprot.org/citations/19759395), PubMed:[28867292](http://www.uniprot.org/citations/28867292), PubMed:[9705271](http://www.uniprot.org/citations/9705271)). The complex (1) mediates the end resection of damaged DNA, which generates proper single-stranded DNA, a key initial steps in HR, and is (2) required for the recruitment of other repair factors and efficient activation of ATM and ATR upon DNA damage (PubMed:[19759395](http://www.uniprot.org/citations/19759395), PubMed:[9705271](http://www.uniprot.org/citations/9705271)). The MRN complex possesses single-strand endonuclease activity and double-strand-specific 3'-5' exonuclease activity, which are provided by MRE11, to initiate end resection, which is required for single-strand invasion and recombination (PubMed:[19759395](http://www.uniprot.org/citations/19759395), PubMed:[28867292](http://www.uniprot.org/citations/28867292), PubMed:[9705271](http://www.uniprot.org/citations/9705271)). Within the MRN complex, NBN acts as a protein-protein adapter, which specifically recognizes and binds phosphorylated proteins, promoting their recruitment to DNA damage sites (PubMed:[12419185](http://www.uniprot.org/citations/12419185), PubMed:[15616588](http://www.uniprot.org/citations/15616588), PubMed:[18411307](http://www.uniprot.org/citations/18411307), PubMed:[18582474](http://www.uniprot.org/citations/18582474), PubMed:[18583988](http://www.uniprot.org/citations/18583988), PubMed:[18678890](http://www.uniprot.org/citations/18678890), PubMed:[19759395](http://www.uniprot.org/citations/19759395), PubMed:[19804756](http://www.uniprot.org/citations/19804756), PubMed:[23762398](http://www.uniprot.org/citations/23762398), PubMed:[24534091](http://www.uniprot.org/citations/24534091), PubMed:[27814491](http://www.uniprot.org/citations/27814491), PubMed:[27889449](http://www.uniprot.org/citations/27889449), PubMed:[33836577](http://www.uniprot.org/citations/33836577)). Recruits MRE11 and RAD50 components of the MRN complex to DSBs in response to DNA damage (PubMed:[12419185](http://www.uniprot.org/citations/12419185), PubMed:[18411307](http://www.uniprot.org/citations/18411307), PubMed:[18583988](http://www.uniprot.org/citations/18583988), PubMed:[18678890](http://www.uniprot.org/citations/18678890), PubMed:[24534091](http://www.uniprot.org/citations/24534091), PubMed:[26438602](http://www.uniprot.org/citations/26438602)). Promotes the recruitment of PI3/PI4-kinase family members ATM, ATR, and probably DNA-PKcs to the DNA damage sites, activating their functions (PubMed:[15064416](http://www.uniprot.org/citations/15064416), PubMed:[15616588](http://www.uniprot.org/citations/15616588), PubMed:[15790808](http://www.uniprot.org/citations/15790808), PubMed:[16622404](http://www.uniprot.org/citations/16622404), PubMed:[22464731](http://www.uniprot.org/citations/22464731), PubMed:[30952868](http://www.uniprot.org/citations/30952868), PubMed:[35076389](http://www.uniprot.org/citations/35076389)). Mediates the recruitment of phosphorylated RBBP8/CtIP to DSBs, leading to cooperation between the MRN complex and RBBP8/CtIP to initiate end resection (PubMed:[19759395](http://www.uniprot.org/citations/19759395), PubMed:[19759395](http://www.uniprot.org/citations/19759395), PubMed:[19759395](http://www.uniprot.org/citations/19759395)).

[27814491](http://www.uniprot.org/citations/27814491) </a>, PubMed: <a href="http://www.uniprot.org/citations/27889449" target="\_blank">27889449 </a>, PubMed: <a href="http://www.uniprot.org/citations/33836577" target="\_blank">33836577 </a>). RBBP8/CtIP specifically promotes the endonuclease activity of the MRN complex to clear DNA ends containing protein adducts (PubMed: <a href="http://www.uniprot.org/citations/27814491" target="\_blank">27814491 </a>, PubMed: <a href="http://www.uniprot.org/citations/27889449" target="\_blank">27889449 </a>, PubMed: <a href="http://www.uniprot.org/citations/30787182" target="\_blank">30787182 </a>, PubMed: <a href="http://www.uniprot.org/citations/33836577" target="\_blank">33836577 </a>). The MRN complex is also required for the processing of R-loops (PubMed: <a href="http://www.uniprot.org/citations/31537797" target="\_blank">31537797 </a>). NBN also functions in telomere length maintenance via its interaction with TERF2: interaction with TERF2 during G1 phase preventing recruitment of DCLRE1B/Apollo to telomeres (PubMed: <a href="http://www.uniprot.org/citations/10888888" target="\_blank">10888888 </a>, PubMed: <a href="http://www.uniprot.org/citations/28216226" target="\_blank">28216226 </a>). NBN also promotes DNA repair choice at dysfunctional telomeres: NBN phosphorylation by CK2 promotes non-homologous end joining repair at telomeres, while unphosphorylated NBN promotes microhomology-mediated end-joining (MMEJ) repair (PubMed: <a href="http://www.uniprot.org/citations/28216226" target="\_blank">28216226 </a>). Enhances AKT1 phosphorylation possibly by association with the mTORC2 complex (PubMed: <a href="http://www.uniprot.org/citations/23762398" target="\_blank">23762398 </a>).

### Cellular Location

Nucleus. Chromosome. Nucleus, PML body. Chromosome, telomere Note=Localizes to discrete nuclear foci after treatment with genotoxic agents (PubMed:10783165, PubMed:26215093, PubMed:26438602). Localizes to DNA double-strand breaks (DSBs); recruited to DNA damage sites via association with phosphorylated proteins, such as phosphorylated H2AX, phosphorylated MDC1 and phosphorylated RAD17 (PubMed:12419185, PubMed:18411307, PubMed:18582474, PubMed:18583988, PubMed:18678890, PubMed:19338747, PubMed:23115235, PubMed:24534091, PubMed:26438602) Acetylation of 'Lys-5' of histone H2AX (H2AXK5ac) promotes NBN/NBS1 assembly at the sites of DNA damage (PubMed:26438602)

### Tissue Location


Ubiquitous (PubMed:9590180). Expressed at high levels in testis (PubMed:9590180).


### Phospho-p95/NBS1 (S343) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Phospho-p95/NBS1 (S343) Antibody - Images





Western blot analysis of p95/NBS1 phosphorylation expression in Jurkat cell lysate treated with Etoposide.