

# c-Jun Antibody

Rabbit mAb Catalog # AP90332

# **Specification**

# c-Jun Antibody - Product Information

Application WB, IHC, ICC, IP

Primary Accession P05412
Reactivity Rat

Clonality Monoclonal

**Other Names** 

AH119; AP-1; AP1; Activator protein 1; Jun A; Proto-oncogene c-jun; RJG-9; V-jun avian sarcoma

virus 17 oncogene homolog; c-Jun; p39;

Isotype Rabbit IgG
Host Rabbit
Calculated MW 35676 Da

# c-Jun Antibody - Additional Information

Dilution WB~~1:1000

IHC~~1:100~500

ICC~~N/A IP~~N/A

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human

c-Jun

Description c-jun is a transcription factor that

recognizes and binds to the enhancer heptamer motif 5'-TGA[CG]TCA-3'. Promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation.

Storage Condition and Buffer

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide

and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid

freeze / thaw cycle.

# c-Jun Antibody - Protein Information

### Name JUN

#### **Function**

Transcription factor that recognizes and binds to the AP-1 consensus motif 5'-TGA[GC]TCA-3' (PubMed:<a href="http://www.uniprot.org/citations/10995748" target="\_blank">10995748</a>, PubMed:<a href="http://www.uniprot.org/citations/22083952" target="\_blank">22083952</a>). Heterodimerizes with proteins of the FOS family to form an AP-1 transcription complex, thereby



enhancing its DNA binding activity to the AP-1 consensus sequence 5'-TGA[GC]TCA-3' and enhancing its transcriptional activity (By similarity). Together with FOSB, plays a role in activation-induced cell death of T cells by binding to the AP-1 promoter site of FASLG/CD95L, and inducing its transcription in response to activation of the TCR/CD3 signaling pathway (PubMed:<a href="http://www.uniprot.org/citations/12618758" target="\_blank">12618758</a>). Promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation (PubMed:<a href="http://www.uniprot.org/citations/17210646" target="\_blank">17210646</a>). Involved in activated KRAS-mediated transcriptional activation of USP28 in colorectal cancer (CRC) cells (PubMed:<a href="http://www.uniprot.org/citations/24623306" target="\_blank">24623306</a>). Binds to the USP28 promoter in colorectal cancer (CRC) cells (PubMed:<a href="http://www.uniprot.org/citations/24623306" target="\_blank">24623306</a>).

**Cellular Location**Nucleus.

#### **Tissue Location**


Expressed in the developing and adult prostate and prostate cancer cells.

# c-Jun Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## c-Jun Antibody - Images



Western blot analysis of c-jun expression in NIH/3T3 cell lysates.