

Tubulin gamma Antibody
Rabbit mAb
Catalog # AP90437**Specification**

Tubulin gamma Antibody - Product Information

Application	WB, IHC, FC, ICC, IP
Primary Accession	P23258
Reactivity	Rat, Dog, Fish
Clonality	Monoclonal

Other Names

Gamma-1-tubulin; Gamma-tubulin complex component 1; GCP-1; TBG1; TUBG; TUBG1; TUBGCP1; Tubulin gamma-1 chain; tubulin, gamma 1; tubulin, gamma polypeptide, gamma tubulin;

Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	51170 Da

Tubulin gamma Antibody - Additional Information

Dilution	WB~~1:1000 IHC~~1:100~500 FC~~1:10~50 ICC~~N/A IP~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Tubulin gamma
Description	TUBG1 Tubulin is the major constituent of microtubules. Gamma tubulin is found at microtubule organizing centers (MTOC) such as the spindle poles or the centrosome. Pericentriolar matrix component that regulates alpha/beta tubulin minus-end nucleation, centrosome duplication and spindle formation. Interacts with GCP2 and GCP3. Interacts with B9D2. Interacts with CDK5RAP2; Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage Condition and Buffer	

Tubulin gamma Antibody - Protein Information

Name TUBG1 ([HGNC:12417](#))

Synonyms TUBG

Function

Tubulin is the major constituent of microtubules, protein filaments consisting of alpha- and beta-tubulin heterodimers (PubMed:38305685, PubMed:38609661, PubMed:39321809). Gamma-tubulin is a key component of the gamma-tubulin ring complex (gTuRC) which mediates microtubule nucleation (PubMed:38305685, PubMed:38609661, PubMed:39321809). The gTuRC regulates the minus-end nucleation of alpha-beta tubulin heterodimers that grow into microtubule protofilaments, a critical step in centrosome duplication and spindle formation (PubMed:38305685, PubMed:38609661, PubMed:39321809).

Cellular Location


Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle. Note=Localizes to mitotic spindle microtubules.

Tubulin gamma Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Tubulin gamma Antibody - Images



Western blot analysis of Tubulin gamma expression in (1) HeLa cell lysate; (2) NIH 3T3 cell lysate; (3) C6 cell lysate; (4) Zebrafish lysates; (5) COS-1 cell lysate; (6) MDCK cell lysate.