

Histone H2B (mono methyl R79) Antibody
Rabbit mAb
Catalog # AP90488**Specification**

Histone H2B (mono methyl R79) Antibody - Product Information

Application	WB
Primary Accession	Q16778
Clonality	Monoclonal
Other Names	
H2B; H2BQ; GL105; H2B.1; H2BFQ; H2BGL105; H2R79me1;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	13920 Da

Histone H2B (mono methyl R79) Antibody - Additional Information

Dilution	WB~~1:1000
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Histone H2B (mono methyl R79)
Description	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a member of the histone H2B family, and generates two transcripts through the use of the conserved stem-loop termination motif, and the polyA addition motif.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Histone H2B (mono methyl R79) Antibody - Protein Information**Name** H2BC21 ([HGNC:4760](#))**Function**

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Cellular Location


Nucleus. Chromosome.

Histone H2B (mono methyl R79) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Histone H2B (mono methyl R79) Antibody - Images



Western blot analysis of Histone H2B (mono methyl R79) expression in HeLa cell lysate.