

Raptor Antibody
Rabbit mAb
Catalog # AP90602**Specification**

Raptor Antibody - Product Information

Application	WB, IHC, FC, ICC, IP
Primary Accession	Q8N122
Reactivity	Rat
Clonality	Monoclonal
Other Names	
RPTOR; KOG1; KIAA1303; RAPTOR; Mip1;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	149038 Da

Raptor Antibody - Additional Information

Dilution	WB~~1:1000 IHC~~1:100~500 FC~~1:10~50 ICC~~N/A IP~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Raptor
Description	The regulatory associated protein of mTOR (Raptor) was identified as an mTOR binding partner that mediates mTOR signaling to downstream targets. Raptor binds to mTOR substrates, including 4E-BP1 and p70 S6 kinase, through their TOR signaling (TOS) motifs and is required for mTOR-mediated phosphorylation of these substrates. Binding of the FKBP12-rapamycin complex to mTOR inhibits the mTOR-raptor interaction, suggesting a mechanism for rapamycin's specific inhibition of mTOR signaling. This mTOR-raptor interaction and its regulation by nutrients and/or rapamycin is dependent on a protein called GβL.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Raptor Antibody - Protein Information

Name RPTOR ([HGNC:30287](#))**Function**

Component of the mechanistic target of rapamycin complex 1 (mTORC1), an evolutionarily conserved central nutrient sensor that stimulates anabolic reactions and macromolecule biosynthesis to promote cellular biomass generation and growth (PubMed:[12150925](http://www.uniprot.org/citations/12150925), PubMed:[12150926](http://www.uniprot.org/citations/12150926), PubMed:[12747827](http://www.uniprot.org/citations/12747827), PubMed:[24403073](http://www.uniprot.org/citations/24403073), PubMed:[26588989](http://www.uniprot.org/citations/26588989), PubMed:[32561715](http://www.uniprot.org/citations/32561715), PubMed:[37541260](http://www.uniprot.org/citations/37541260)). In response to nutrients, growth factors or amino acids, mTORC1 is recruited to the lysosome membrane and promotes protein, lipid and nucleotide synthesis by phosphorylating several substrates, such as ribosomal protein S6 kinase (RPS6KB1 and RPS6KB2) and EIF4EBP1 (4E-BP1) (PubMed:[12150925](http://www.uniprot.org/citations/12150925), PubMed:[12150926](http://www.uniprot.org/citations/12150926), PubMed:[12747827](http://www.uniprot.org/citations/12747827), PubMed:[24403073](http://www.uniprot.org/citations/24403073), PubMed:[26588989](http://www.uniprot.org/citations/26588989), PubMed:[37541260](http://www.uniprot.org/citations/37541260)). In the same time, it inhibits catabolic pathways by phosphorylating the autophagy initiation components ULK1 and ATG13, as well as transcription factor TFEB, a master regulators of lysosomal biogenesis and autophagy (PubMed:[12150925](http://www.uniprot.org/citations/12150925), PubMed:[12150926](http://www.uniprot.org/citations/12150926), PubMed:[12747827](http://www.uniprot.org/citations/12747827), PubMed:[24403073](http://www.uniprot.org/citations/24403073), PubMed:[32561715](http://www.uniprot.org/citations/32561715), PubMed:[37541260](http://www.uniprot.org/citations/37541260)). The mTORC1 complex is inhibited in response to starvation and amino acid depletion (PubMed:[12150925](http://www.uniprot.org/citations/12150925), PubMed:[12150926](http://www.uniprot.org/citations/12150926), PubMed:[12747827](http://www.uniprot.org/citations/12747827), PubMed:[24403073](http://www.uniprot.org/citations/24403073), PubMed:[37541260](http://www.uniprot.org/citations/37541260)). Within the mTORC1 complex, RPTOR acts both as a molecular adapter, which (1) mediates recruitment of mTORC1 to lysosomal membranes via interaction with small GTPases Rag (RagA/RRAGA, RagB/RRAGB, RagC/RRAGC and/or RagD/RRAGD), and a (2) substrate-specific adapter, which promotes substrate specificity by binding to TOS motif-containing proteins and direct them towards the active site of the MTOR kinase domain for phosphorylation (PubMed:[12747827](http://www.uniprot.org/citations/12747827), PubMed:[24403073](http://www.uniprot.org/citations/24403073), PubMed:[26588989](http://www.uniprot.org/citations/26588989), PubMed:[37541260](http://www.uniprot.org/citations/37541260)). mTORC1 complex regulates many cellular processes, such as odontoblast and osteoclast differentiation or neuronal transmission (By similarity). mTORC1 complex in excitatory neuronal transmission is required for the prosocial behavior induced by the psychoactive substance lysergic acid diethylamide (LSD) (By similarity).

Cellular Location

Lysosome membrane. Cytoplasm Cytoplasmic granule. Note=Targeting to lysosomes depends on amino acid availability: recruited to lysosome membranes via interaction with GTP-bound form of RagA/RRAGA (or RagB/RRAGB) in complex with the GDP-bound form of RagC/RRAGC (or RagD/RRAGD), promoting recruitment of mTORC1 to the lysosomes (PubMed:31601708, PubMed:31601764). In arsenite-stressed cells, accumulates in stress granules when associated

with SPAG5 and association with lysosomes is drastically decreased (PubMed:23953116).

Tissue Location


Highly expressed in skeletal muscle, and in a lesser extent in brain, lung, small intestine, kidney and placenta

Raptor Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Raptor Antibody - Images



Western blot analysis of Raptor expression in MCF-7 cell lysate.