

RALBP1 Antibody

Rabbit mAb Catalog # AP90679

Specification

RALBP1 Antibody - Product Information

Application Primary Accession Reactivity Clonality Other Names RALBP1; RIP1; RLIP1; RLIP76;	WB, IHC, FC <u>015311</u> Rat Monoclonal
lsotype Host Calculated MW	Rabbit IgG Rabbit 76063 Da
RALBP1 Antibody - Additional Information	
Dilution	WB~~1:1000

Dilution	WB~~1:1000
	IHC~~1:100~500
	FC~~1:10~50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human RALBP1
Description	RalBP1 interacts with RalA and the endocytosis protein REPS2 (POB1) through its carboxy-terminal Ral binding domain. RalBP1 has an intrinsic GTPase activating function and interacts with Cdc42 through its centrally located Rho-GAP domain. A protein complex containing RalBP1/POB1/RalA regulates endocytosis of membrane receptors. RalBP1 also functions as a non-ABC transporter that catalyzes the ATP-dependent transport of numerous xenobiotics, including glutathione conjugates and some
Storage Condition and Buffer	chemotherapeutic agents. Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short
	term. Store at -20°C long term. Avoid freeze / thaw cycle.

RALBP1 Antibody - Protein Information

Name RALBP1 (HGNC:9841)



Function

Multifunctional protein that functions as a downstream effector of RALA and RALB (PubMed:7673236). As a GTPase-activating protein/GAP can inactivate CDC42 and RAC1 by stimulating their GTPase activity (PubMed: 7673236). As part of the Ral signaling pathway, may also regulate ligand-dependent EGF and insulin receptors-mediated endocytosis (PubMed:10910768, PubMed:12775724). During mitosis, may act as a scaffold protein in the phosphorylation of EPSIN/EPN1 by the mitotic kinase cyclin B-CDK1, preventing endocytosis during that phase of the cell cycle (PubMed:12775724). During mitosis, also controls mitochondrial fission as an effector of RALA (PubMed:21822277). Recruited to mitochondrion by RALA, acts as a scaffold to foster the mitotic kinase cyclin B-CDK1-mediated phosphorylation and activation of DNM1L (PubMed: 21822277).

Cellular Location

Cell membrane; Peripheral membrane protein. Cytoplasm, cytosol Cytoplasm, cytoskeleton, spindle pole {ECO:000250|UniProtKB:Q62796} Nucleus. Mitochondrion. Note=Cytosolic protein that transiently associates with the mitotic spindle poles in early prophase, and dissociates from them after completion of mitosis (By similarity) Targeted to the plasma membrane through its interaction with RALB, directed by FGF signaling. Docking on the membrane is required to transduce the Ral signal (By similarity). Recruited by RALA to the mitochondrion during mitosis where it regulates mitochondrial fission (PubMed:21822277). Nuclear localization is cell cycle dependent while membrane localization is seen in adherent cells (PubMed:22319010). The region involved in membrane association could form transmembrane domains and expose a part of the protein extracellularly (Probable) {ECO:000250|UniProtKB:Q62796, ECO:000250|UniProtKB:Q9PT60, ECO:0000269|PubMed:21822277, ECO:000269|PubMed:22319010, ECO:0000305|PubMed:15610018}

Tissue Location

Expressed ubiquitously but at low levels. Shows a strong expression in the erythrocytes.


RALBP1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

RALBP1 Antibody - Images





Western blot analysis of RALBP1 expression in HeLa cell lysate.