

**Moesin Antibody**  
**Rabbit mAb**  
**Catalog # AP90724****Specification**

---

**Moesin Antibody - Product Information**

Application	WB, IHC, FC, ICC, IP
Primary Accession	<a href="#">P26038</a>
Reactivity	Rat
Clonality	Monoclonal
<b>Other Names</b>	
MSN; Moesin;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	67820 Da

**Moesin Antibody - Additional Information**

Dilution	WB~~1:1000 IHC~~1:100~500 FC~~1:10~50 ICC~~N/A IP~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Moesin
Description	The ezrin, radixin, and moesin (ERM) proteins function as linkers between the plasma membrane and the actin cytoskeleton and are involved in cell adhesion, membrane ruffling, and microvilli formation. ERM proteins undergo intra or intermolecular interaction between their amino- and carboxy-terminal domains, existing as inactive cytosolic monomers or dimers.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

**Moesin Antibody - Protein Information****Name** MSN ([HGNC:7373](#))**Function**

Ezrin-radixin-moesin (ERM) family protein that connects the actin cytoskeleton to the plasma

membrane and thereby regulates the structure and function of specific domains of the cell cortex. Tethers actin filaments by oscillating between a resting and an activated state providing transient interactions between moesin and the actin cytoskeleton (PubMed:<a href="http://www.uniprot.org/citations/10212266" target="\_blank">10212266</a>). Once phosphorylated on its C-terminal threonine, moesin is activated leading to interaction with F-actin and cytoskeletal rearrangement (PubMed:<a href="http://www.uniprot.org/citations/10212266" target="\_blank">10212266</a>). These rearrangements regulate many cellular processes, including cell shape determination, membrane transport, and signal transduction (PubMed:<a href="http://www.uniprot.org/citations/12387735" target="\_blank">12387735</a>, PubMed:<a href="http://www.uniprot.org/citations/15039356" target="\_blank">15039356</a>). The role of moesin is particularly important in immunity acting on both T and B-cells homeostasis and self-tolerance, regulating lymphocyte egress from lymphoid organs (PubMed:<a href="http://www.uniprot.org/citations/9298994" target="\_blank">9298994</a>, PubMed:<a href="http://www.uniprot.org/citations/9616160" target="\_blank">9616160</a>). Modulates phagolysosomal biogenesis in macrophages (By similarity). Also participates in immunologic synapse formation (PubMed:<a href="http://www.uniprot.org/citations/27405666" target="\_blank">27405666</a>).

### Cellular Location

Cell membrane; Peripheral membrane protein {ECO:0000250|UniProtKB:P26041}; Cytoplasmic side {ECO:0000250|UniProtKB:P26041}. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:P26041}. Apical cell membrane {ECO:0000250|UniProtKB:P26041}; Peripheral membrane protein {ECO:0000250|UniProtKB:P26041}; Cytoplasmic side {ECO:0000250|UniProtKB:P26041}. Cell projection, microvillus membrane {ECO:0000250|UniProtKB:P26041}; Peripheral membrane protein {ECO:0000250|UniProtKB:P26041}; Cytoplasmic side {ECO:0000250|UniProtKB:P26041}. Cell projection, microvillus {ECO:0000250|UniProtKB:P26041}. Note=Phosphorylated form is enriched in microvilli-like structures at apical membrane. Increased cell membrane localization of both phosphorylated and non-phosphorylated forms seen after thrombin treatment (By similarity). Localizes at the uropods of T lymphoblasts. {ECO:0000250|UniProtKB:P26041, ECO:0000269|PubMed:18586956, ECO:0000269|PubMed:9298994}

### Tissue Location


In all tissues and cultured cells studied.


### Moesin Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Moesin Antibody - Images





Western blot analysis of Moesin expression in HeLa cell lysate.